



## Investigation of solid-liquid phase diagrams of the sulfamethazine-salicylic acid co-crystal

Dipali Ahuja, Michael Svärd, Åke C. Rasmuson

### Publication date

01-01-2019

### Published in

CrystEngComm;21 (18), pp. 2837-2972

### Licence

This work is made available under the [CC BY-NC-SA 1.0](#) licence and should only be used in accordance with that licence. For more information on the specific terms, consult the repository record for this item.

### Document Version

1

### Citation for this work (HarvardUL)

Ahuja, D., Svärd, M. and Rasmuson, Å.C. (2019) 'Investigation of solid-liquid phase diagrams of the sulfamethazine-salicylic acid co-crystal', available: <https://hdl.handle.net/10344/8913> [accessed 24 Jul 2022].

This work was downloaded from the University of Limerick research repository.

For more information on this work, the University of Limerick research repository or to report an issue, you can contact the repository administrators at [ir@ul.ie](mailto:ir@ul.ie). If you feel that this work breaches copyright, please provide details and we will remove access to the work immediately while we investigate your claim.

## **Supporting Information**

### **Investigation of solid-liquid phase diagrams of the Sulfamethazine-Salicylic acid Co-crystal**


Dipali Ahuja,<sup>a</sup> Michael Svärd,<sup>a</sup> and Åke C. Rasmuson<sup>a</sup>

<sup>a</sup>*Synthesis and Solid State Pharmaceutical Centre, Bernal Institute, Department of Chemical Sciences, University of Limerick, Co. Limerick, Ireland*


## **Table of Contents**

<b><u>Title</u></b>	<b><u>Figure/ Table</u></b>
Full (top) and zoom-in view (bottom) of ternary phase diagram of SMT-SA co-crystal system in methanol at 10°C.	S1
Full (top) and zoom-in view (bottom) of ternary phase diagram of SMT-SA co-crystal system in methanol at 20°C.	S2
Ternary phase diagram of SMT-SA co-crystal system in methanol at 30°C.	S3
Full (top) and zoom-in view (bottom) of ternary phase diagram of SMT-SA co-crystal system in acetonitrile at 10°C.	S4
Full (top) and zoom-in view (bottom) of ternary phase diagram of SMT-SA co-crystal system in acetonitrile at 10°C.	S5
Ternary phase diagram of SMT-SA co-crystal system in acetonitrile at 30°C.	S6
Ternary phase diagram of SMT-SA co-crystal system in 7:3 (v/v) DMSO-methanol mixture at 20°C.	S7
Ternary phase diagram of SMT-SA co-crystal system in 7:3 (v/v) DMSO-methanol mixture at 40°C.	S8
A zoom-in view of the ternary phase diagram of SMT-SA co-crystal system in acetonitrile depicting the solubility points and invariant points at 10 (red), 20 (blue) and 30°C (green) showing the effect of temperature.	S9
A zoom-in view of the ternary phase diagram of SMT-SA co-crystal system in methanol depicting the solubility points and invariant points at 10 (red), 20 (blue) and 30°C (green) showing the effect of temperature.	S10
A zoom-in view of the ternary phase diagram of SMT-SA co-crystal system in 7:3 (v/v) DMSO-methanol mixture depicting the solubility points and invariant points at 20 (red), 30 (blue) and 40°C (green) showing the effect of temperature.	S11
PXRD patterns of the pure SMT, 1:1 SMT-SA co-crystal and the invariant points at 10 (red), 20 (blue) and 30°C (green) in methanol where SMT and co-crystal co-exist.	S12
DSC thermograms of the invariant points at 10 (red), 20 (blue) and 30°C (green) in methanol where SMT and co-crystal co-exist.	S13
PXRD patterns of the pure SA, 1:1 SMT-SA co-crystal and the invariant points at 10 (red), 20 (blue) and 30°C (green) in methanol where SA and co-crystal co-exist.	S14
DSC thermograms of the invariant points at 10 (red), 20 (blue) and 30°C	S15


(green) in methanol where SA and co-crystal co-exist.	
PXRD patterns of the pure SMT, 1:1 SMT-SA co-crystal and the invariant points at 10 (red), 20 (blue) and 30°C (green) in acetonitrile where SMT and co-crystal co-exist.	S16
DSC thermograms of the invariant points at 10 (red), 20 (blue) and 30°C (green) in acetonitrile where SMT and co-crystal co-exist.	S17
PXRD patterns of the pure SA, 1:1 SMT-SA co-crystal and the invariant points at 10 (red), 20 (blue) and 30°C (green) in acetonitrile where SA and co-crystal co-exist.	S18
DSC thermograms of the invariant points at 10 (red), 20 (blue) and 30°C (green) in acetonitrile where SA and co-crystal co-exist.	S19
PXRD patterns of the pure SMT, 1:1 SMT-SA co-crystal and the invariant points at 10 (red), 20 (blue) and 30°C (green) in 7:3 (v/v) DMSO-methanol mixture where SMT and co-crystal co-exist.	S20
DSC thermograms of the invariant points at 10 (red), 20 (blue) and 30°C (green) in 7:3 (v/v) DMSO-methanol mixture where SMT and co-crystal co-exist.	S21
PXRD patterns of the pure SA, 1:1 SMT-SA co-crystal and the invariant points at 10 (red), 20 (blue) and 30°C (green) in 7:3 (v/v) DMSO-methanol mixture where SA and co-crystal co-exist.	S22
DSC thermograms of the invariant points at 10 (red), 20 (blue) and 30°C (green) in 7:3 (v/v) DMSO-methanol mixture where SA and co-crystal co-exist.	S23




**Figure S1.** Full (top) and zoom-in view (bottom) of ternary phase diagram of SMT-SA co-crystal system in methanol at 10°C. The points, regions and line explained in Figure 4 of article.



**Figure S2.** Full (top) and zoom-in view (bottom) of ternary phase diagram of SMT-SA co-crystal system in methanol at 20°C. The points, regions and line explained in Figure 4 of article.




**Figure S3.** Ternary phase diagram of SMT-SA co-crystal system in methanol at 30°C. The points, regions and line explained in Figure 4 of article.




**Figure S4.** Full (top) and zoom-in view (bottom) of ternary phase diagram of SMT-SA co-crystal system in acetonitrile at 10°C. The points, regions and line explained in Figure 4 of article.






**Figure S5.** Full (top) and zoom-in view (bottom) of ternary phase diagram of SMT-SA co-crystal system in acetonitrile at 10°C. The points, regions and line explained in Figure 4 of article.







**Figure S6.** Ternary phase diagram of SMT-SA co-crystal system in acetonitrile at 30°C. The points, regions and line explained in Figure 4 of article.




**Figure S7.** Ternary phase diagram of SMT-SA co-crystal system in 7:3 (v/v) DMSO-methanol mixture at 20°C. The points, regions and line explained in Figure 4 of article.




**Figure S8.** Ternary phase diagram of SMT-SA co-crystal system in 7:3 (v/v) DMSO-methanol mixture at 40°C. The points, regions and line explained in Figure 4 of article.




**Figure S9.** A zoom-in view of the ternary phase diagram of SMT-SA co-crystal system in acetonitrile depicting the solubility points and invariant points at 10 (red), 20 (blue) and 30°C (green) showing the effect of temperature.




**Figure S10.** A zoom-in view of the ternary phase diagram of SMT-SA co-crystal system in methanol depicting the solubility points and invariant points at 10 (red), 20 (blue) and 30°C (green) showing the effect of temperature.




**Figure S11.** A zoom-in view of the ternary phase diagram of SMT-SA co-crystal system in 7:3 (v/v) DMSO-methanol mixture depicting the solubility points and invariant points at 20 (red), 30 (blue) and 40°C (green) showing the effect of temperature.




**Figure S12.** PXRD patterns of the pure SMT, 1:1 SMT-SA co-crystal and the invariant points at 10 (red), 20 (blue) and 30°C (green) in methanol where SMT and co-crystal co-exist.



**Figure S13.** DSC thermograms of the invariant points at 10 (red), 20 (blue) and 30°C (green) in methanol where SMT and co-crystal co-exist.




**Figure S14.** PXRD patterns of the pure SA, 1:1 SMT-SA co-crystal and the invariant points at 10 (red), 20 (blue) and 30°C (green) in methanol where SA and co-crystal co-exist.




**Figure S15.** DSC thermograms of the invariant points at 10 (red), 20 (blue) and 30°C (green) in methanol where SA and co-crystal co-exist.







**Figure S16.** PXRD patterns of the pure SMT, 1:1 SMT-SA co-crystal and the invariant points at 10 (red), 20 (blue) and 30°C (green) in ACETONITRILE where SMT and co-crystal co-exist.




**Figure S17.** DSC thermograms of the invariant points at 10 (red), 20 (blue) and 30°C (green) in acetonitrile where SMT and co-crystal co-exist.




**Figure S18.** PXRD patterns of the pure SA, 1:1 SMT-SA co-crystal and the invariant points at 10 (red), 20 (blue) and 30°C (green) in acetonitrile where SA and co-crystal co-exist.




**Figure S19.** DSC thermograms of the invariant points at 10 (red), 20 (blue) and 30°C (green) in acetonitrile where SA and co-crystal co-exist.




**Figure S20.** PXRD patterns of the pure SMT, 1:1 SMT-SA co-crystal and the invariant points at 10 (red), 20 (blue) and 30°C (green) in 7:3 (v/v) DMSO-methanol mixture where SMT and co-crystal co-exist.



**Figure S21.** DSC thermograms of the invariant points at 10 (red), 20 (blue) and 30°C (green) in 7:3 (v/v) DMSO-methanol mixture where SMT and co-crystal co-exist.



**Figure S22.** PXRD patterns of the pure SA, 1:1 SMT-SA co-crystal and the invariant points at 10 (red), 20 (blue) and 30°C (green) in 7:3 (v/v) DMSO-methanol mixture where SA and co-crystal co-exist.



**Figure S23.** DSC thermograms of the invariant points at 10 (red), 20 (blue) and 30°C (green) in 7:3 (v/v) DMSO-methanol mixture where SA and co-crystal co-exist.