



## Pillow project data set

Maebh Barry

### Publication date

22-03-2018

### Licence

This work is made available under the **CC BY-NC-SA 1.0** licence and should only be used in accordance with that licence. For more information on the specific terms, consult the repository record for this item.

### Document Version

1

### Citation for this work (HarvardUL)

Barry, M. (2018) 'Pillow project data set', available: <https://hdl.handle.net/10344/6675> [accessed 25 Jul 2022].

This work was downloaded from the University of Limerick research repository.

For more information on this work, the University of Limerick research repository or to report an issue, you can contact the repository administrators at [ir@ul.ie](mailto:ir@ul.ie). If you feel that this work breaches copyright, please provide details and we will remove access to the work immediately while we investigate your claim.

## Correct answers Infant CPR and actions if infant choking

Initial Actions	Number correct (N=44)	Percentage correct
1. Checks for responsiveness	29	65.91
2. Initial Call for help if helper there or speaker mobile	25	56.82
3. Remove any obstruction if possible	32	73
<b>Step 1 Puff</b>		
Places one hand on the forehead - finger under chin tilt to open airway	35	79.55
Makes a seal around nose and mouth with your mouth	36	81.82
Blows gently and steadily for one second	36	81.82
Looks for infant chest rise	26	59.09
Indicates 5 puffs or rescue breaths	34	77.27
<b>Step 2 Pump or chest compressions</b>		
Places two fingers in centre of baby's chest	43	97.73
Compresses 1/3 of the chests depth (Depth of Pillow)	40	90.91
Releases pressure and allow the chest to come back up	38	86.36
Indicates rate of 30 compressions to two puffs	42	95.45
Indicates 100 to 120 times per minute (quick)	39	88.64
Indicates after one minute to call 999 /112	41	93.18
Continue 30 pumps and 2 puffs until help arrives	43	97.73
<b>Initial Actions infant Choking</b>		
Check for obvious obstruction	39	88.64
Supports head face down lower than the body and gives five back slaps	40	90.91
Checks the mouth and removes object if seen with finger and thumb only	41	93.18
Places two fingers nipple line push in and upwards five times	37	84.09
Check the mouth for the object	34	77.27
After three cycles of back slaps and chest thrusts call for an ambulance	38	86.36
Continues with chest thrusts and back slaps until help arrives	38	86.36