

ULRR

The effect of strength training on performance in endurance athletes.

Item Type	Article
Authors	Beattie, Kris;Kenny, Ian C.;Lyons, Mark;Carson, Brian P.
Citation	Sports Medicine 44 (6), pp. 845-65
Publisher	Springer
Download date	2026-05-09 14:13:17
Item License	https://creativecommons.org/licenses/by-nc-sa/1.0/
Link to Item	https://hdl.handle.net/10344/4182

THE EFFECT OF STRENGTH TRAINING ON PERFORMANCE IN ENDURANCE ATHLETES

Authors: Kris Beattie, Ian C. Kenny, Mark Lyons and Brian P. Carson.
Title: **The Effect of Strength Training on Performance in Endurance Athletes**
Address: Department of Physical Education and Sport Sciences, University of Limerick,
Limerick, Ireland
Contact Details: Email: kris.beattie@ul.ie
Telephone: 00353 61 234781
Fax: 00353 61 202814

THE EFFECT OF STRENGTH TRAINING ON PERFORMANCE IN ENDURANCE ATHLETES

ABSTRACT

BACKGROUND

Economy, velocity/power at maximal oxygen uptake ($v \dot{V}O_{2 \max} / w \dot{V}O_{2 \max}$) and endurance-specific muscle power tests (i.e. maximal anaerobic running velocity v_{MART}), are now thought to be the best performance predictors in elite endurance athletes. In addition to cardiovascular function, these key performance indicators are believed to be partly dictated by the neuromuscular system. One technique to improve neuromuscular efficiency in athletes is through strength training.

OBJECTIVE

The aim of this systematic review was to search the body of scientific literature for original research investigating the effect of strength training on performance indicators in well-trained endurance athletes - specifically economy, $v \dot{V}O_{2 \max} / w \dot{V}O_{2 \max}$ and muscle power (v_{MART}).

METHODS

A search was performed using MEDLINE, PubMed, ScienceDirect, SPORTDiscus and Web of Science search engines. There were twenty-six studies that met the inclusion criteria (athletes had to be trained endurance athletes with ≥ 6 months endurance training, training ≥ 6 hours per week OR $\dot{V}O_{2 \max} \geq 50$ ml/min/kg, the strength interventions had to be ≥ 5 weeks in duration, and control groups used). All studies were reviewed using the PEDro scale.

RESULTS

The results showed that strength training improved time trial performance, economy, $v \dot{V}O_{2 \max} / w \dot{V}O_{2 \max}$ and v_{MART} in competitive endurance athletes.

CONCLUSION

The present research available supports the addition of strength training in an endurance athlete's programme for improved economy, $v \dot{V}O_{2 \max} / w \dot{V}O_{2 \max}$, muscle power and performance. However, it is evident that further research is needed. Future investigations should include valid strength assessments (i.e. squats, jump-squats, drop jumps) through a range of velocities (maximal strength \leftrightarrow strength-speed \leftrightarrow speed-strength \leftrightarrow reactive-strength), and administer appropriate strength programmes (exercise, load & velocity prescription) over a long-term intervention period (> 6 months) for optimal transfer to performance.

1. INTRODUCTION

Endurance sport performance relies on a complex inter-play of physiological and biomechanical factors. Cardiovascular capacity has often been thought to be the main limiting factor in endurance performance. Classical measures such as maximal oxygen uptake ($\dot{V}O_{2\max}$) and lactate threshold (LT) have been traditionally used in the laboratory to predict the performance potential of runners, cyclists, triathletes and cross-country skiers.^[1] Consequently, physical preparation for these sports has generally focused on developing these two physiological qualities. However, elite endurance athletes with similar $\dot{V}O_{2\max}$ levels can have differing abilities during a race and therefore maximum oxygen uptake cannot fully explain true racing ability. Economy, and assessments that include an endurance-specific muscle power component such as velocity/power during maximal oxygen uptake ($v\dot{V}O_{2\max} / w\dot{V}O_{2\max}$) and maximal anaerobic running velocity (v MART), are now thought to be superior performance indicators in an elite population.^[2]

Economy is the amount of metabolic energy expended at a given velocity or power output.^[3] Economical movement is multi-factorial and is determined by training history, anthropometrics, biomechanics and physiology.^[4] During a race, an economical athlete will use less energy at sub-maximal intensities and spare vital carbohydrate stores for significant stages in competition (i.e. sprint finish). East Africans have dominated distance running for the past few decades and it is believed that their success is partly due to their superior running economy.^[3] Improvements in economy may be difficult to obtain in highly-trained endurance athletes and therefore any novel training modality that results in marginal improvements may be crucial for success.

Endurance-specific muscle power is the ability of the neuromuscular system to rapidly produce force following a sustained period of high-intensity exercise (high glycolytic and/or oxidative energy demand)^[5]. This ability may be the differentiating factor for elite endurance performance as successful athletes at world-level can produce high velocities and power outputs to win a race following a sustained period of high-intensity exercise (i.e. sprint finish). Therefore rate of force development (RFD) is essential not only in sprint and power sports, but also in elite endurance competition. Endurance-specific muscle power assessments such as peak velocity during the maximal anaerobic running test (v MART) have been found to be better predictors of running performance in an elite population because they are both highly influenced by neuromuscular and anaerobic factors.^[2] The v MART consists of a series of incremental 20 second runs with 100 second recoveries on a treadmill until volitional exhaustion.^[6] Peak velocity/power at $\dot{V}O_{2\max}$ ($v\dot{V}O_{2\max}$), is influenced by $\dot{V}O_{2\max}$, economy and lactate threshold. However is also shown to have a large 'muscle power' component because it is strongly correlated to v MART ($r = 0.85$, $p < 0.001$).^[2] McLaughlin et al.^[7] found that in well-trained runners $v\dot{V}O_{2\max}$ was the best predictor of running performance over 16 km. Also, Millet et al.^[8] found that peak power output during an incremental cycling test (W_{peak}) was correlated to overall performance in elite triathletes. Consequently, in addition to cardiovascular ability, limitations to elite endurance performance may be dictated by other dynamical system factors, including neuromuscular function.

THE EFFECT OF STRENGTH TRAINING ON PERFORMANCE IN ENDURANCE ATHLETES

One training technique for improving muscle force-velocity function in athletes is through strength training.^[9] It is proposed that through neuromuscular adaptations (musculotendinous stiffness, motor unit recruitment and synchronisation, rate coding, intra- and inter-muscular coordination, and neural inhibition) strength training has the potential to improve performance in endurance athletes through increased (1) economy, and (2) endurance-specific muscle power factors (i.e. v MART).^[2] Theoretically, a strength-trained endurance athlete will (1) be more economical as sub-maximal forces developed during each stride or pedal revolution would decrease to a lower percentage of maximal values, and (2) have improved endurance-specific muscle power as they are able to produce higher maximum running or cycling velocities through an improved ability to rapidly absorb and create force against the ground or pedal (Figure 1).




Fig 1. Hypothetical model of the determinants for elite endurance performance and the potential benefits from strength training (LSD = long slow distance training; intervals = repeated bouts of exercise lasting ~1 to 8 minutes and eliciting an oxygen demand equal to ~90 to 100% of $\dot{V}O_{2max}$; PCr = phosphocreatine; $\dot{V}O_{2max}$ = maximal O₂ uptake; v MART = peak velocity in maximal anaerobic running test; $v\dot{V}O_{2max}$ = peak velocity at $\dot{V}O_{2max}$). The red font and bold arrows highlight the potential benefit of strength training on endurance performance [Adapted from Paavolainen et al.^[5] with permission].

THE EFFECT OF STRENGTH TRAINING ON PERFORMANCE IN ENDURANCE ATHLETES

Elite endurance athletes are renowned for their high volume of (low force) endurance training. Unfortunately, unlike strength training, specific endurance training such as ‘interval’ or ‘tempo’ sessions are not effective in improving neuromuscular function in well-trained endurance athletes (Figure 1). Traditionally, for unknown reasons, endurance athletes have been cautious to strength train. In fact, research investigating the training characteristics of runners competing in the 2008 U.S. Olympic Marathon trials found that they “*included little strength training in their training programs...and nearly half the runners did no strength training at all.*”^[10] This philosophy may be due to endurance athletes and coaches being uneducated in strength training science and the associated potential performance improvements. The aim of this systematic review was to search the body of scientific literature for original research investigating the effect of strength training on performance, specifically economy and assessments that include an endurance-specific muscle power component (i.e. $\dot{V}O_{2\max} / w$, $\dot{V}O_{2\max}$, and $\dot{V}MART$), in well-trained endurance athletes.

THE EFFECT OF STRENGTH TRAINING ON PERFORMANCE IN ENDURANCE ATHLETES

2. METHODS

A search was performed using MEDLINE, PubMed, ScienceDirect, SPORTDiscus and Web of Science search engines to identify studies that assessed the effect of strength training on performance in competitive endurance athletes. The following keywords were used in the search (“strength training” OR “resistance training” OR “weight training” OR “weightlifting” OR “concurrent training” OR “plyometrics”) AND (“endurance athletes” OR “cyclists” OR “runners” OR “triathletes” OR “cross-country skiers”) AND (“performance”). Strength training was defined as non-cycling/running/cross-country skiing, weight-loaded activity including bodyweight, free-weight and machine-based exercises. The sub-categories for strength training included: (1) maximal-strength training that targets maximal force development through high-load, low-velocity movements (i.e. squats, deadlifts), (2) explosive-strength training (strength-speed & speed-strength) that improves rate of force development (RFD) and maximal power output through medium- to high-load, high-velocity movements (i.e. squat jumps, Olympic lifts); and (3) reactive-strength training that targets musculotendinous stiffness and stretch-shortening cycle function through low-load, high-velocity exercises (i.e. jumps, drop-jumps, hops, bounds, sprints).

Inclusion criterion for this analysis were (1) athletes had to be trained endurance athletes (≥ 6 months endurance training, training ≥ 6 hours per week, $\dot{V}O_{2max} \geq 50$ ml/min/kg), (2) the strength interventions had to be ≥ 5 weeks in duration, and (3) control groups had to be used. All articles were read and the outcomes of each study summarised. Articles were excluded if the study methodology did not meet the specific inclusion criteria. Other relevant articles were obtained through additional bibliographical means (Figure 2).




Fig 2. PRISMA (Preferred Reporting Items for Systematic Reviews) flow chart illustrating the inclusion and exclusion criteria used in the systematic review. PEDro indicates Physiotherapy Evidence Database.

THE EFFECT OF STRENGTH TRAINING ON PERFORMANCE IN ENDURANCE ATHLETES

The Physiotherapy Evidence Database (PEDro) scale was used to rate the quality of the selected articles. The PEDro scale is an 11-item scale designed for rating the methodological quality of randomised controlled trials (Maher et al 2003). Each satisfied item (except for the first item, which relates to external validity) contributes 1 point to the total PEDro score.^[11] The items include random allocation; concealment of allocation; comparability of groups at baseline; blinding of subjects, researchers, and assessors; analysis by intention to treat; and adequacy of follow up. The PEDro scale ranges from 0 to 10, where 0 points (the worst possible score) are awarded to a study that fails to satisfy any of the included items and 10 points (the best possible score) are awarded to a study that satisfies all the included items. Studies scoring 9 or 10 on the PEDro scale are considered to have methodologically excellent internal validity, those scoring 6 to 8 are considered good, those scoring 4 or 5 are fair, and those scoring less than 4 are poor. All studies graded using the PEDro scale were included.

3. RESULTS

Twenty six papers met the inclusion criteria. Of these papers, eight were from running, nine from cycling, six from cross-country skiing and three from triathlon. Tables 1 – 3 compare the results. The tables are subdivided into the four sports (running, cycling, cross-country skiing and triathlon) and are structured to compare (i) subjects (sample size, sex, standard of racing, $\dot{V}O_{2\max}$, weekly training volume) and research design (PEDro score, group allocation, control of training) [Table 1], (ii) strength intervention (type of strength training, programme overview, frequency and duration of training [Table 2] and (iii) results [Table 3].

3.1 PEDro score analysis

Scores on the PEDro scale for the twenty six selected articles ranged from 5 to 6 of a maximum 10 points. Only fourteen studies randomly allocated their subjects into training groups and scored 6 out of 10 on the PEDro scale. [12-25] The additional twelve studies scored 5 out of 10: four studies did not mention randomised allocation of subjects [26-29] and four studies allowed the subjects to select their own groups. [30-33] Other studies allocated subjects into training groups by $\dot{V}O_{2\max}$, [34] $\dot{V}O_{2\max}$ and 5 km time trial performance, [5] mean training time; [35] or by randomly allocating half of the subjects into groups and then the rest by age and 5 km time trial performance. [36]

3.2 Running (time trial performance, $\dot{V}O_{2\max}$ and economy)

In runners, improvements were found in time trial performance, economy, $\dot{V}O_{2\max}$ and vMART after a strength training intervention. The studies show that 8 weeks of explosive-strength training can improve 3 km time trial performance [15], and reactive-strength training can significantly improve 5 km [5] ($p < 0.05$) and 3 km [13] ($p < 0.05$, ES = 0.13) performance. Both Mikkola et al. [27] and Berryman et al. [15] both found an increase in $\dot{V}O_{2\max}$ from 8 weeks of both reactive-strength and explosive-strength training. The two studies that assessed vMART both found a significant ($p < 0.01$) improvement following an 8 week [27] and 9 week [5] reactive-strength programme. Five studies found significant improvements in economy from both maximal- [12, 36] and reactive-strength training interventions. [5, 13, 15]

3.3 Cycling (time trial performance, $\dot{V}O_{2\max}$ and economy)

In cyclists, 12-16 weeks of maximal-strength training was found to significantly improve 5 minute [30] ($p < 0.01$) and 45 minute time trial performance [19] ($p < 0.05$, ES = 0.66). Improvements were also found in 40 minute [31] and 60 minute time trial ability; [35] however these improvements were not found to be significantly different to their allocated control groups. From the six cycling studies that analysed power at $\dot{V}O_{2\max}$ ($\dot{V}O_{2\max}$), three found improvements [28, 30, 35] but only Rønnestad et al.'s work [28, 30] found a significant effect when compared against the control group ($p < 0.05$, ES = 0.81 [28], ES = 84 [30]). Bastiaans et al [35] found significant improvements in 'delta efficiency' ($p < 0.05$, ES = 0.49), and Rønnestad et al [30] showed increases in economy and 'work efficiency' during the final 60 minutes of a 185 minute cycle test ($p < 0.05$).

3.4 Cross-country skiing (time trial performance and economy)

THE EFFECT OF STRENGTH TRAINING ON PERFORMANCE IN ENDURANCE ATHLETES

In cross-country skiers, Losnegard *et al* ^[33] found a significant increase in a 1.1 km ‘upper body double-poling’ time trial ($p < 0.05$), as well as a non-significant improvement in a 1.3 km ‘full-body roller ski’ time trial from their strength training intervention. Mikkola *et al* ^[26] also found a significant improvement in 2 km ‘upper-body double-poling,’ however there was no significant difference in change between the control and the experimental group. Rønnestad *et al* ^[32] found no improvement in 7.5 km ‘full-body roller ski’ time trial performances. Improvements in economy were seen for both ‘whole-body roller skiing’ ^[32] ($p < 0.05$, ES = 0.77) and ‘isolated upper-body double-poling’ movements ^[21, 22, 26].

3.5 Triathlon ($\dot{V}O_2 \max_v$, $\dot{V}O_2 \max_w$ and economy)

In triathletes, Millet *et al.* ^[23] found a significant increase in peak treadmill velocity at $\dot{V}O_2 \max$ ($p < 0.01$, ES = 0.55) following a maximal-strength training intervention, whereas Hausswirth *et al.* ^[24] found no difference in $\dot{V}O_2 \max_w$ during a cycling protocol. Out of the three studies that investigated running economy in triathletes, only Millet *et al* ^[23] found significant increases at 25% ($p < 0.05$, ES = 1.15) and 75% $\dot{V}O_2$ ($p < 0.05$, ES = 0.14).

THE EFFECT OF STRENGTH TRAINING ON PERFORMANCE IN ENDURANCE ATHLETES

SYSTEMATIC REVIEW

THE EFFECT OF STRENGTH TRAINING ON PERFORMANCE IN ENDURANCE ATHLETES

Table 1. Studies included in the meta-analysis: subjects and research design											
Reference	Subjects					Research design					
	n	Sex	Age (y)	VO _{2max} (mL/min/kg)	Level; weekly volume/hours; duration of competitiveness	PE德罗 score	Assigned to group?	Intervention (n)	Control (n)	ET controlled?	ST replacement or addition?
Running											
Johnston et al. ^[12]	12	F	30.3	50.5	32-48 km/ week for > 1 year	6	RCT	6	6	Yes	Addition
Paavolainen et al. ^[5]	18	M	23	67.7	Elite cross-country	5	Matched with regard to VO ₂ and 5 km TT	10	8	Yes	Replacement
Spurrs et al. ^[13]	17	M	25	57.6	Trained; 60-80 km/week for 10 years	6	RCT	8	9	Yes - monitored	Addition
Saunders et al. ^[14]	15	M	23.4	71.1	6 internationals, all national; 107 km/week	6	RCT	7	8	Yes - training duration matched	Addition
Mikkola et al. ^[27]	25	M and F	17	62.1	Post-pubertal, high school runners	5	No mention of RCT	13	12	Yes -volume	Replacement
Støren et al. ^[36]	17	M and F	29.2	59.9	Trained	5	Half RCT, other half matched for 5km and age	8	9	Yes – volume & intensity	Addition
Berryman et al. ^[15]	28	M	28	56.9	Provincial standard, 3-5 sessions per week	6	RCT	11 (reactive) 12 (explosive)	5	Yes – volume & intensity	Addition
Fletcher et al. ^[16]	12	M	24.3	67.5	Highly trained; 70-170 km/week	6	RCT	6	6	-	-
Cycling											
Bastiaans et al. ^[35]	14	M	25	-	6 ± 6 years competing	5	Matched for mean training time	6	8	Yes – HR and training zones	Replacement
Jackson et al. ^[17]	23	18 M, 5 F	30	52	≥0.5 years competing	6	RCT	High Res 9, High Rep 9	5	Yes – HR and training zones	Addition
Levin et al. ^[18]	14	M	31	62.75	≥1 years competing	6	RCT	7	7	Monitored but not controlled	Addition
Rønnestad et al. ^[30]	20	18 M, 2 F	28.5	66.35	Well-trained	5	Self-chosen	11	9	Yes – HR and training zones	Addition
Rønnestad et	20	18 M, 2	28.5	66.35	Norwegian national-	5	Self-chosen	11	9	Yes – HR and	Addition

THE EFFECT OF STRENGTH TRAINING ON PERFORMANCE IN ENDURANCE ATHLETES

al. ^[31]		F			level					training zones	
Sunde et al. ^[34]	13	10 M, 3 F	32.85	61.05	Well-trained and competitive	5	Matched for $\dot{V}O_{2max}$	8	5	Yes – HR and training zones	Addition
Rønnestad et al. ^[28]	12	11 M, 1 F	30	66.25	Norwegian national-level	5	-	6	6	Yes – HR and training zones	Addition
Aagaard et al. ^[19]	14	M	19.5	72.5	U23 international	6	RCT	7	7	Yes – HR and training zones	Addition
Rønnestad et al. ^[29]	27	25 M, 2 F	27.6	63.4	20 highly-trained, 7 recreational	5	-	11 (cyclists) 9 (recreational)	7 (cyclists)	Yes – HR and training zones	Replacement
Cross-country skiing											
Hoff et al. ^[20]	15	F	17.9	55.3	8.8 h / week	6	RCT	8	7	Yes	Replacement of strength-endurance
Hoff et al. ^[21]	19	M	19.8	69.4	'well-trained'	6	RCT	9	10	Yes	Replacement of strength-endurance
Osteras et al. ^[22]	19	M	22.7	61.2	'highly trained' > 5 years	6	RCT	10	9	Yes	Replacement of strength-endurance
Mikkola et al. ^[26]	19	M	23.1	66.5	Finnish national (6-15 years)	5	-	8	11	-	Replacement
Rønnestad et al. ^[32]	17	M	19.5	66.2	National and international Nordic combined	5	No – self selected	8	9	Yes	Replacement
Losnegard et al. ^[33]	19	M and F	21.5	64.7	National	5	No – self selected	9	10	-	-
Triathlon											
Millet et al. ^[23]	15	-	22.85	68.7	20.4 h/week; elite/international	6	RCT	7	8	Yes - 'recorded'	Addition
Hauswirth et al. ^[24]	14	M	31.3	69.2	17.3 h/week; regional and national level	6	RCT	7	7	Yes – 'strictly aerobic , 75% HR'	Addition
Bonnacci et al. ^[25]	8	M and F	21.6	-	Competed for 4.4 years	6	RCT	3	5	No	Addition

Values are means except where stated otherwise (Abbreviations: $\dot{V}O_{2max}$ = maximal oxygen uptake; PEDro score = physiotherapy evidence database score; ET = endurance training; ST = strength training; HR = heart rate; h = hours; M = male; F = female).

THE EFFECT OF STRENGTH TRAINING ON PERFORMANCE IN ENDURANCE ATHLETES

Table 2. Studies included in the meta-analysis: strength interventions						
Reference	Type	Programme overview/example	Closed-chain leg exercises?	Frequency	Duration (wk)	Time of year
Running						
Johnston et al. ^[12]	Maximal-strength	3 x 6RM (parallel squat, seated press, hammer curl, lung, heel-raise and bench press) 3 x 8RM (knee flexion/extension, lateral pull down and seated row) 2 x 20RM (bent leg heel-raise), 2 x 12RM (straight leg heel-raise) and 2 x 15RM weighted sit-up	Yes – squat and lunge	3 x week	10	-
Paavolainen et al. ^[5]	Reactive-strength	Sprints and jumps Alternative jumps, bilateral countermovement, drop and hurdle jumps, 1-legged, 5 jumps	Yes – all reactive exercises	-	9	Off-season
Spurrs et al. ^[13]	Reactive-strength	W 1 60 contacts, W2 100, W3 136, W4 150, W5 170, W6 180. Plyo progression: Squat Jump, split scissor jump, double leg bound, SL hops, depth jump, DL hurdle hop, SL hurdle hop	Yes – all reactive exercises	W 1-3: 2 x week, W 4-6: 3 x week	6	-
Saunders et al. ^[14]	Reactive-strength	Session 1 (Back extension, leg press, CMJs, knee lifts, ankle jumps, hamstring curls) Session 2 (bounds, skips, SL ankles, hurdle jumps, scissors for height)	Yes – all reactive exercises	3 x week	9	-
Mikkola et al. ^[27]	Reactive-strength	Sprints (5-10 x 30-150m), pogos, squat jumps, half squats, knee extensions, calf raises, curls (2-3 x 6-10 reps)	Yes – all reactive exercises	3 x week	8	Pre-competition
Støren et al. ^[36]	Maximal-strength	4 x 4 half squats	Yes – squats	3 x week	8	-
Berryman et al. ^[15]	Reactive- and explosive-strength	Reactive group – drop jumps Explosive group – concentric squat jumps P _{max} load	Yes – drop jumps & concentric squats	1 x week	8	-
Fletcher et al. ^[16]	Maximal- / isometric-strength	4 x 20s at 80% MVC isometric plantar flexion	No – isolated isometric plantar flexion	3 x week	8	Pre-competition
Cycling						

THE EFFECT OF STRENGTH TRAINING ON PERFORMANCE IN ENDURANCE ATHLETES

Bastiaans et al. ^[35]	Muscular endurance	4 x 30 (squats, leg press, step-up) and 2 x 30 (leg pull and core)	Yes – squats and Smith machine step-ups	3 x week	9	Pre-season
Jackson et al. ^[17]	Muscular endurance and maximal-strength	Wk 1 – all 2 x 20, Wk 2 – 10 High Res (4 x 4RM), High Rep (2 x 20RM) ALL squats, leg press, leg curl, Smith machine step-ups, planks	Yes – squats and Smith machine step-ups	3 x week	10	In-season
Levin et al. ^[18]	Maximal-strength, explosive-strength & muscular endurance	Strength 4 x 5 (lunges, squats, RDLs, calf raises crunches) Power 3 x 6 (Jumps squats, SL jump squats, clean grip deadlift, calf raise back extension) Endurance 3 x 12 (SL leg press, knee extension, knee flexion, calf raise & crunches)	Yes – squats, lunges, RDLs, deadlifts etc	3 x week	6	Pre-season
Rønnestad et al. ^[30]	Maximal-strength	W 1-3: 10RM Session 1, 6RM Session 2 W 4-6: 8 RM & 5RM W 7-12: 6RM & 4RM ALL half-squat smith, SL leg press, hip flexion & toe raise.	Yes – Smith squat	2 x week	12	Pre-season
Rønnestad et al. ^[31]	Maximal-strength	W 1-3: 10RM Session 1, 6RM Session 2 W 4-6: 8 RM & 5RM W 7-12: 6RM & 4RM All half-squat smith, SL leg press, hip flexion & toe raise.	Yes – Smith squat	2 x week	12	Pre-season
Sunde et al. ^[34]	Maximal-strength	4 x 4RM half-squats (Smith machine)	Yes – Smith squat	3 x week	8	Pre-season
Rønnestad et al. ^[28]	Maximal-strength	W 1-3: 10RM Session 1, 6RM Session 2 W 4-6: 8 RM & 5RM W 7-12: 6RM & 4RM ALL half-squat Smith, SL leg press, hip flexion & toe raise. W13-25 (SEASON): 2 x 5 (half squat & leg press) 1 x 6 (hip flexion & ankle plantar flexion)	Yes – Smith squat	2 x week	25	Pre-season prep (12W) & In-season (12W)
Aagaard et al. ^[19]	Maximal-strength	W1: 3 x 12, W2-3: 3 x 10, W4-5: 3 x 8, W6-16: 2-	No – all machine isolated	2-3 x week	16	-

THE EFFECT OF STRENGTH TRAINING ON PERFORMANCE IN ENDURANCE ATHLETES

		3x6 (knee extension, leg press, hamstring curl & calf raises)				
Rønnestad et al. ^[29]	Maximal-strength	W 1-3: 10RM Session 1, 6RM Session 2 W 4-6: 8 RM & 5RM W 7-12: 6RM & 4RM ALL half-squat smith, SL leg press, hip flexion & toe raise.	Yes – Smith squat	2 x week	12	-
Levin et al. ^[18]	Maximal-strength, explosive-strength & muscular endurance	Strength 4 x 5 (lunges, squats, RDLs, calf raises crunches) Power 3 x 6 (Jumps squats, SL jump squats, clean grip deadlift, calf raise back extension) Endurance 3 x 12 (SL leg press, knee extension, knee flexion, calf raise & crunches)	Yes – Squats, lunges, RDLs, deadlifts	3 x week	6	Pre-season
Cross-country skiing						
Hoff et al. ^[20]	Maximal-strength	Pull-downs – 3 x 6 Increased by 1kg every session (control group used their normal 'strength-endurance' programme <60% 1RM)	No	3 x week	9	Pre-season
Hoff et al. ^[21]	Maximal-strength	Pull-downs – 3 x 6 Increased by 3kg every session (control group used their normal 'strength-endurance' programme <85% 1RM)	No	45 min / week	8	Pre-season
Østerås et al. ^[22]	Maximal-strength	Pull-downs – 3 x 6 Increased by 3kg every session (control group used their normal 'strength endurance' programme <85% 1RM)	No	45min / week	9	Pre-season
Mikkola et al. ^[26]	Explosive- & reactive-strength	Day 1: Specific explosive - double poling sprints 10 x 10 seconds Day 2: General explosive – half squat, pull over, leg	Yes	3 x week	8	-

THE EFFECT OF STRENGTH TRAINING ON PERFORMANCE IN ENDURANCE ATHLETES

		press, lat pull-down 3 x 6-10 Day 3: Reactive – running sprints, jumps, skating jumps, pogos 3-6 x 20m				
Rønnestad et al. ^[32]	Maximal-strength	Deep squat: W1-6 (3-5x4-8), W7-12 (4-5x3-5) Seated pull-down: W1-6 (3x6-10), W7-12 (3x5-8) Standing double poling	Yes	2 x week	12	-
Losnegard et al. ^[33]	Maximal-strength	Half-squat, pull-down, seated pull-down, double poling, triceps press. W1-3 (3x6-10), W4 (3x5-8), W5-8 (4x8), W9-12 (3x4-6)	Yes	2 x week (W1-8) 1 x week (W9-12)	12	Pre-season
Triathlon						
Millet et al. ^[23]	Maximal-strength	W1 3 x 5, W2 4 x 5, W3 5 x 5 Hamstring curl, leg press, seated press, parallel squat, leg extension and heel-raise	Yes – parallel squat	2 x week	14	Pre-season
Hauswirth et al. ^[24]	Maximal-strength	3-5 x 3-5 Leg press, leg extension, hamstring curl, calf raise.	No	3 x week	5	Pre-season
Bonnacci et al. ^[25]	Reactive-strength	CMJs, knee lifts, pogos, squats, bounds, skips, scissors etc.	Yes	3 x week	8	-

Values are means except where stated otherwise (Abbreviations: CMJ = countermovement jump; W = week; HR = heart rate; h = hours; M = male; F = female; RM = repetition maximum; RDLs = Romanian deadlifts; SL = single-leg).

THE EFFECT OF STRENGTH TRAINING ON PERFORMANCE IN ENDURANCE ATHLETES

Table 3. Studies included in the meta-analysis: results									
Reference	Tests	Strength	Economy	vVO _{2max} ^b	vMART	TT	PP	TTE	Body composition/other performance
Running									
Johnston et al. ^[12]	Squat, knee flexion, body composition, RE and VO _{2max}	↑ Squat (40%), knee flexion (27%) [p < 0.05] ^a	[mL/kg/min] ↑ at 214 m/min (4%) [ES = 0.72] and 230 m/min (ES = 0.64) [p < 0.05] ^a	-	-	-	-	-	Increased body mass and fat-free mass (NS)
Paavolainen et al. ^[5]	5 km TT, isometric knee extension, VO _{2max} , LT, RE, vMART, v20m, 5BJ	↑ Isometric MVC, v20m, 5BJ (p < 0.01)	[mL/kg/min] ↑ at 4.17 m/s (8.1%) [p < 0.001]	- (used gradient)	↑ (p < 0.01) ^a	5 km (3.1%) [p < 0.05]	-	-	Increased body mass, calf and thigh girth (NS)
Spurrs et al. ^[13]	RE, VO _{2max} , LT, MTS, Isometric MVC, RFD, CMJ, 5BJ, 3km TT	↑ Isometric MVC (12.5%) MTS @ 75% MVC (12.9%) RFD (14.5%) CMJ (13.2%) 5BJ (7.8%) (p < 0.05)	[ml/kg/min] ↑ at 12km/h (7.7%, ES = 0.45), 14 km/h (6.4%, ES = 0.45) & 16km/h (4.1%, ES = 0.3) (p < 0.05) ^a	- (used gradient)	-	↑3 km (2.7%, ES = 0.13) ^a	-	-	Increased in body mass NS
Saunders et al. ^[14]	RE, VO _{2max} , 5CMJ, RFD	↑ 5CMJ (15%), RFD (14%) NS	[L/min] ↑ at 18km/h (4.1%) (p = 0.02, ES = 0.35) ^a but NS at 14km/h, 16km/h	-	-	-	-	-	Increased in body mass NS
Mikkola et al. ^[27]	ISO MVC, vMART, RE, 30m, 5J, CMJ, VO _{2max} , v VO _{2max}	↑ Isometric MVC (8%), 1RM leg extension (4%), RFD (31%) (p < 0.05) ^a . No sig changes in CMJ & 5J	[ml/kg/min] ↑ at 12, 13 & 14km/h NS	↑ 1.2% NS	↑ 3% (p < 0.01)	-	-	-	Increased lean body mass, calf & thigh girth NS ↑V30m (1.1%) (p < 0.05)
Støren et al. ^[36]	1RM half squat, RFD, RE, TTE at MAS,	↑* 1RM (33.2%), RFD (26%) of half squat (p < 0.01)	[ml/kg ^{0.75} /min] ↑ at 70% VO _{2max} (5%) (p < 0.01, ES = 1.03) ^a	- (used gradient)	-	-	-	↑ TTE at MAS (21.3%) (p < 0.05) ^a	Increased body mass NS
Berryman et al. ^[15]	VO _{2max} , vVO ₂ , economy, P _{peak} , 3 km TT, RE	↑ P _{peak} (W) in both reactive & explosive group	[ml/kg ^{0.75} /min] ↑ in both reactive (ES = 0.96)	↑ in both reactive (ES = 0.49) &	-	↑3km TT In reactive (ES = 0.46)	-	-	No changes in body mass

THE EFFECT OF STRENGTH TRAINING ON PERFORMANCE IN ENDURANCE ATHLETES

		(p < 0.01)	& explosive (ES = 0.63) groups (p < 0.01)	explosive (ES = 0.43) groups (p < 0.01)		& explosive (ES = 0.37) (p < 0.05)			
Fletcher et al. ^[16]	VO _{2max} , LT, RE, isometric triceps surae tendon stiffness (TST)	No improvement ISO TST	[kJ/kg/km] No improvement	- (used gradient)	-	-	-	-	
Cycling									
Bastiaans et al. ^[35]	60 min TT, incremental W _{max} , DE and 30s power	-	[delta efficiency] ↑ 1.41% (p < 0.05, ES = 0.49)	↑ W _{max} 4.7% (p < 0.01, ES = 0.64) ^b		[60 min TT] ↑ 7.9% (p < 0.01, ES = 0.86) ^b	-	-	All groups ↑ (NS)
Jackson et al. ^[17]	Squats, leg curls, SL press, step-ups, VO _{2max}	-	- (only examined peak economy at VO _{2max})	NS		No mean change over 30 km test	-	-	-
Levin et al. ^[18]	30 km test (with 250m & 1 km power), 1RM squat & VO _{2max}	↑ 1RM squat NS	-	↓		No mean change over 30 km Test	No sig diff except for last 1 km sprint of 30 km test (p < 0.05, ES 0.3)		Increased body mass NS
Rønnestad et al. ^[30]	1RM half-squat smith, VO _{2max} , 185min at 44% W _{max} + 5min TT	↑ 1RM half smith-squat (26%) (p < 0.01)	[ml/kg/min] ↑ (p < 0.05) economy during 185 min at 44% W _{max} , ↑ during final 60 min (p < 0.05) ^a	↑ W _{max} (4.2%) (p < 0.05) ^a		[5min TT] ↑ power 7% (p < 0.01)	-		Increased body mass NS Increased knee flexors/extensors CSA
Rønnestad et al. ^[31]	Muscle CSA, isometric half-squat, VO _{2max} , Wingate, 40min TT	↑ isometric strength (21.2%) (p < 0.01)	-	↑ W _{max} (4.3%) (p < 0.05, ES = 0.44)		[40min TT] ↑ power (6%) (p < 0.01, ES = 0.57) ^a	[30s Wingate PP] ↑ (9.4%) PP (p < 0.01, ES 0.61)		↑ TTE at maximal aerobic power (p < 0.05, ES 0.57) ^a Increase body mass NS
Sunde et al. ^[34]	Smith squat 1RM, RFD, CE @ 70%, TTE, VO _{2max} , LT	↑ 1RM Smith Squat (14.2%) ↑ RFD Smith squat (16.2%) (p < 0.05)	↑ [WE] (4.7%) (p < 0.05, ES = 0.48) ^a and [ml/kg ^{0.67} /W] ↑ (3%) at 70% VO _{2max} (p < 0.05, ES = 0.56)	-		-	-		Increased knee extensor/flexor CSA (p < 0.05)
Rønnestad et al. ^[28]	Muscle CSA, Half-squat, VO _{2max} , Wingate, 40min TT	↑ 1RM half-smith squat (23%) 12 weeks and was maintained to 25 weeks (p < 0.01)	-	↑ W _{max} (8%) (p < 0.05, ES = 0.81)		-	[Wingate PP] ↑ PP (p < 0.05, ES = 0.67) ^a		No change in muscle CSA
Aagaard et al. ^[19]	Isometric knee	↑ isometric	[ml/Joule]	-		[45min TT]			Reduce freely chosen

THE EFFECT OF STRENGTH TRAINING ON PERFORMANCE IN ENDURANCE ATHLETES

	extensor MVC, RFD, 45 min TT	MVC (12%) (p < 0.05) & RFD (20%) (p < 0.01)	No change in strength group			↑ 8% (p < 0.01, ES = 0.66)			cadence Increased patellar tendon CSA
Rønnestad et al. ^[29]	VO _{2max} , 1RM smith half squat, 5min at 125W for VO ₂	↑1RM smith-squat (31%) (p < 0.01)	[ml/kg/min] ↑ economy at 125W but NS	-		-			All groups ↑ (NS)
Cross-country skiing									
Hoff et al. ^[20]	1 RM DP pull-down, peak force and RFD at 80% and 60% 1RM, VO _{2max} (running), VO _{2peak} upper body 'poling', TTE upper, economy at max	↑ 1RM, time to peak force at 80% 1RM (p < 0.05)	[UB DP (mL/kg ^{0.67} /m)] ↑(p < 0.001) ^b	- (UB DP VO _{2max})	-	-	-	↑ TTE (p < 0.001)	
Hoff et al. ^[21]	1 RM DP pull-down, peak force and RFD @ 80% and 60% 1RM, VO _{2max} (running), VO _{2peak} upper body 'poling', TTE upper, economy at max	↑ 1RM (9.9%), peak force at 80% (34%) and 60% (33%) 1RM (p < 0.05)	[UB DP (mL/kg ^{0.67} /m)] ↑1t 1.81 m/min (p < 0.05)	- (UB DP VO _{2max})	-	-	-	↑ TTE (56%) at V _{O2} peak velocity (p < 0.05)	
Østerås et al. ^[22]	1RM 'ski pull-down' F-V, P-V at various loads VO _{2peak} , TTE	↑ power & velocities at each load (except lowest) (p < 0.01)	[UB DP (ml/kg ^{0.67} /min)] ↑* double poling at pre-test VO _{2peak} force (p < 0.01, ES = 1.66)	- (UB DP VO _{2max})	-	-	-	↑* TTE at VO _{2max} velocity (p < 0.05, ES = 1.18)	
Mikkola et al. ^[26]	Leg extensor isometric & concentric force-time, 30m double poling with roller skis, Velocity & economy 2km UB double	↑leg extensor ISO & CON NS	[UB DP (ml/kg/min)] ↑* during constant velocity 2km (7%) (p < 0.05)	- (walking VO ₂ max with poles)	-	No change 2km UB poling velocity	-		Increased lean body mass, ↑ 30m (1.4%) double poling, (p < 0.05)

THE EFFECT OF STRENGTH TRAINING ON PERFORMANCE IN ENDURANCE ATHLETES

	poling, VO _{2max} (walking with poles), MAST (maximal anaerobic ski test)								
Rønnestad et al. [32]	1RM squat, pull-down, squat jump height, VO _{2max} roller ski, economy, 7.5km TT	↑ 1RM squat (12%) pull-down (23%), squat jump (8.8%) (p < 0.01)	[Roller Ski (ml/kg/min)] ↑ at 5° (3.8%) (p < 0.05, ES = 0.77) ^a but no change at 4°	-	-	No change rollerski 7.5 km TT	-		Increased vastus lateralis thickness (p < 0.05) No change in body mass
Losnegard et al. [33]	1RM half squat & seated pull-down, CMJ, VO _{2max} rollerski	↑ 1RM half-squat (12%) pull down (19%) (p < 0.01) no change in CMJ.	[Roller Ski (ml/kg/min)] unchanged in both groups	-	-	↑ UB 1.1 km TT (7%) (p < 0.05) but NS rollerksi 1.3 km TT (3.7%)	-		No change in 20m, 40m, 80m & 100m velocity, ↑ 5 min W/kg double-poling (p < 0.05), No change in quadriceps CSA
Triathlon									
Millet et al. [23]	Concentric half squat & heel raise, 10s hopping & limb stiffness Running analysis: VO _{2max} and vVO _{2max} on track, economy at 25% and 75% vVO ₂ during 3km, VO ₂ kinetics	↑ 1RM half squat & heel raise, (p < 0.01) ↑ hopping height and power (p < 0.05) Hopping stiffness not different	[ml/kg/min] ↑ at 25% (ES 1.15) and 75% (ES = 1.14) vVO ₂ during 3 km (p < 0.05)	↑ (p < 0.01, ES = 0.55)	-	-	-	-	No change in VO ₂ kinetics No change in body mass
Hauswirth et al. [24]	1RM leg press, isometric knee extension Cycling analysis: VO _{2max} , PVO _{2max} , gross efficiency	↑ 1RM leg press (6.6%) (p < 0.01) ↑ isometric knee but NS	No differences in gross efficiency	Remain unchanged	-	-	-	-	No change in body mass
Bonnacci et al. [25]	Running analysis: Economy, EMG (for muscle recruitment patterns running	No tests for strength	12km/h NS	-	-	-	-	-	Bike to run testing protocol No change in body mass, thigh or calf girth.

THE EFFECT OF STRENGTH TRAINING ON PERFORMANCE IN ENDURANCE ATHLETES

	after cycling)							
--	----------------	--	--	--	--	--	--	--

Values are means except where stated otherwise. (Abbreviations: ↑ = improved; a = significant difference between strength group pre-and post-test only; b = except for cycling studies, for which the parameter is $\dot{V}O_{2max}$; NS = no significant difference between strength group pre- and post-test; TT = time trial; RE = running economy; LT = lactate threshold; v_{MART} = peak velocity in maximal anaerobic running test; $v_{\dot{V}O_{2max}}$ = peak velocity at $\dot{V}O_{2max}$; $w_{\dot{V}O_{2max}}$ = peak power at $\dot{V}O_{2max}$; MVC = maximum voluntary contraction; RFD = rate of force development; CMJ = countermovement jump; 5 BJ = five broad jump test; ISO = isometric; MAS = maximum aerobic speed; TTE = time to exhaustion; DE = delta economy; SL= single-leg; CSA = cross-sectional area; PP = peak power)

THE EFFECT OF STRENGTH TRAINING ON PERFORMANCE IN ENDURANCE ATHLETES

SYSTEMATIC REVIEW

4. DISCUSSION

Despite the abundance of studies investigating concurrent strength and endurance training, relatively few have examined well-trained endurance athletes. This systematic review is unique due to the focused analysis of strength training on specific performance indicators (economy, $\dot{V}O_{2\max}/W$, $\dot{V}O_{2\max}$, $\dot{V}MART$ and time-trials) in well-trained runners, cyclists, triathletes and cross-country skiers.

4.1 STRENGTH DIAGNOSTICS

As expected, the majority of the reviewed studies demonstrated an improvement in muscle force-velocity characteristics following a strength intervention. [5, 12, 13, 15, 17, 19, 20-22, 27, 28, 31-34, 36] However, it is important to highlight that there were a wide variety of exercises administered throughout the literature to measure maximal-, explosive- and reactive-strength adaptations. Running, cycling, triathlon and cross-country skiing all require the hip, knee and ankle musculature to work in unison to produce force against the ground or pedal. A valid strength test for these sports would measure the force capabilities of the leg extensors in the same way – through closed-chain, multi-joint exercises such as squats, jump-squats or drop jumps. [37] However, some studies in this review [5, 19, 24, 27] assessed strength ability through open-chain, isolated exercises (i.e. knee extension, leg press). Testing force production in an isolated manner may have reduced the validity of the overall force capabilities of the endurance athlete's leg musculature. Another criticism is that most studies only measured force output in one or two velocity ranges, either through low-velocity (one repetition maximum) or high-velocity (unloaded jumps and hops) exercises. It is important to measure force output through a range of velocities to determine maximal-, explosive- (strength-speed & speed-strength) and reactive-strength ability. [38] Assessing force capabilities with valid exercises through a range of velocities would highlight sensitive changes in strength qualities following an intervention period and allow for a more accurate relationship between strength adaptation and endurance performance.

4.1.1 Reactive-strength diagnostics in runners and triathletes

Runners and triathletes need to have proficient eccentric muscular capabilities to rapidly absorb and utilise the elastic energy produced during each ground contact. The short ground contact phase in running is the only phase in which a runner or triathlete can produce force and influence running velocity. Paavolainen et al [5] demonstrated the importance of reactive-strength by finding a strong relationship between ground contact time and running economy ($r = 0.64$, $p < 0.001$). Reactive-strength is affected by musculotendinous stiffness and stretch-shortening cycle (SSC) function. [39] Schmidtbleicher [40] demonstrated that the SSC can be classified as either slow or fast. Fast SSC is characterised by short contact times (< 0.25 seconds) and small angular displacement of the hip, knee and ankle joint; whereas slow SSC involves longer contact times (> 0.25 seconds) and larger angular joint displacements. Unfortunately, the running and triathlon studies in the current review did not take into consideration fast or slow SSC function and only assessed reactive-strength through 'general' reactive-strength measurements such as countermovement jumps, [13, 27] broad jumps and hopping tests. [5, 13, 14, 23] The 'reactive-strength index' (RSI) is a

THE EFFECT OF STRENGTH TRAINING ON PERFORMANCE IN ENDURANCE ATHLETES

popular assessment used by Strength & Conditioning coaches to examine the relationship between force production and ground contact time through a series of drop-jumps at differing heights.^[41] The RSI test may have been a more appropriate and sensitive assessment to track reactive-strength adaptations and transferability to running and triathlon performance.

4.2 TIME TRIAL PERFORMANCE

In well-trained endurance athletes, the current literature indicates that strength training can significantly improve 3 km^[13] ($p < 0.05$, ES = 0.13) and 5 km^[5] ($p < 0.05$) time trial performance in runners, 5 minute^[30] ($p < 0.01$) and 45 minute time trial performance^[19] ($p < 0.05$, ES = 0.66) in cyclists and 1.1 km ‘upper body double-poling’ time trial performance in cross-country skiers ($p < 0.05$). However, it is important to note that elite endurance racing success is not dictated by average velocity or power output over a set distance and therefore time-trial ability is not a ‘true’ reflection of racing performance.^[42] Further analysis of economy and assessments that include an endurance-specific muscle power component (i.e. $\dot{V}O_{2\max} / w$ and $\dot{V}O_{2\max}$, and v MART) may add to the potential beneficial effect of strength training on performance in well-trained endurance athletes.

4.3 ECONOMY

Economy is represented by energy expenditure and is normally expressed as submaximal $\dot{V}O_2$ at a given velocity or power output. It is now established that economy is a critical factor for success in elite endurance sport.^[43] The present research shows that there were significant improvements in economy from both maximal-^[12, 36] and reactive-strength training interventions^[5, 13, 15] in well-trained runners. This supports Noakes^[44] philosophy that runners with poor economy may lack musculotendinous stiffness and therefore strength training may improve the ability of the leg musculature to rapidly absorb and utilise the elastic energy produced during each ground contact. Also in cyclists, the literature shows that strength training significantly improved ‘delta efficiency’^[35] ($p < 0.05$, ES = 0.49), economy during the final 60 minutes of a 185 minute cycle test^[30] ($p < 0.05$) and ‘work efficiency.’^[34] In cross-country skiers, improvements in economy were found in both ‘whole-body roller skiing’^[32] ($p < 0.05$, ES = 0.77) and ‘isolated upper-body double-poling’ movements^[21, 22, 26]. Out of the three studies that investigated running economy in triathletes, only Millet *et al*^[23] found significant increases at 25% ($p < 0.05$, ES = 1.15) and 75% $\dot{V}O_2$ ($p < 0.05$, ES = 0.14).

Interestingly, improvements in economy were found to be velocity specific in runners. Spurrs *et al.*^[13] found a 6.7% and 6.4% significant increase at both 12 km/h (ES = 0.45) and 14 km/h (ES = 0.45), but only a 4.1 % increase at 16 km/h ($p < 0.05$, E = 0.3). Furthermore, Saunders *et al.*^[14] only found a significant improvement at 18 km/h in elite international runners ($p = 0.02$, ES = 0.35), with no change at 14 km/h and 16 km/h. This supports Berg’s^[14] view on adaptation specificity that marathoners may be more economical at marathon pace than 800m and 1500m specialists, whereas middle distance runners may be more efficient at higher velocities. Consequently, the most valid measurement of economy may be at specific race velocities and power outputs, rather than an arbitrary

submaximal intensity which is commonly used. Future researchers should take this into consideration when assigning velocities for economy assessment.

4.4 ENDURANCE MUSCLE POWER

Endurance-specific muscle power is the ability of the neuromuscular system to rapidly produce force following a sustained period of high-intensity exercise (high glycolytic and/or oxidative energy demand) ^[5]. This combined neuromuscular and anaerobic ability may be the differentiating factor for elite endurance performance as successful athletes at world-level can produce high velocities and power outputs to win a race following a sustained period of high-intensity exercise (i.e. sprint finish). As illustrated in Figure 1, $v\dot{V}O_{2\max}$ is not only dictated by $\dot{V}O_{2\max}$, LT and economy, but also by muscle power factors (neuromuscular and anaerobic ability). Noakes ^[47] originally suggested that velocity at $\dot{V}O_{2\max}$ ($v\dot{V}O_{2\max}$) could be used as a potential measure of muscle power in runners. From this review, only Mikkola et al. ^[27] and Berryman et al. ^[15] assessed $v\dot{V}O_{2\max}$. Both researchers found an increase in $v\dot{V}O_{2\max}$ after an 8 week reactive-strength program, with only the latter study showing a significant effect from both reactive- ($p < 0.01$, ES = 0.49) and explosive-strength ($p < 0.01$, ES = 0.43) programmes. From the six cycling studies that analysed power at $\dot{V}O_{2\max}$ ($w\dot{V}O_{2\max}$), three found improvements ^[28, 30, 35] but only Rønnestad et al.'s work ^[28, 30] found a significant effect when compared against the control group ($p < 0.05$, ES = 0.81^[28], ES = 84^[30]). In triathletes, Millet et al. ^[23] established a significant increase in peak treadmill velocity at $\dot{V}O_{2\max}$ ($p < 0.01$, ES = 0.55), whereas Hausswirth et al. ^[24] found no difference in $w\dot{V}O_{2\max}$ during a cycling protocol.

Conversely, Paavolainen et al. ^[2] argues that the aerobic system is still strongly involved during a $\dot{V}O_{2\max}$ test and $v\dot{V}O_{2\max}/w\dot{V}O_{2\max}$ should not be used as a pure measure of endurance-specific muscle power performance. The v MART (peak velocity attained during a maximal anaerobic running test), which consists of a series of incremental 20 seconds sprints on a treadmill until exhaustion, is believed to place more emphasis on assessing neuromuscular and anaerobic performance. The two running studies that assessed v MART in this review both found a significant ($p < 0.01$) improvement following an 8 week ^[27] and 9 week ^[5] reactive-strength programme.

4.5 INTERVENTION ANALYSIS

4.5.1 Programme Duration

Aside from Rønnestad et al.'s ^[28] strength intervention lasting 25 weeks, the average intervention period in this review was approximately 10 weeks. Much of what we know about neurological and structural adaptations in strength training derives from similar short term (8-12 week) interventions involving relatively untrained or inexperienced subjects. ^[48] There are only a few studies investigating the long-term strength adaptations in well-trained athletes; however these are from strength and power sports. ^[49] Future research in well-trained endurance athletes should focus on long-term strength interventions (12-18 months) and subsequent endurance performance.

4.5.2 *Exercise Prescription*

There were a variety of the strength programmes administered all of the twenty-six investigations. The two main distinctions in the interventions are in the prescription of (i) exercises, and (ii) loads and velocities of exercises (see section 4.5.3). ‘Transfer of training’ is a term used to describe the effectiveness of adaptations from a strength exercise transferring to sporting performance.^[50] The ability to generate force is dependent on the limb and joint positioning of the leg extensors.^[51] Therefore, the exercises selected in a program can influence the magnitude of neuromuscular adaptations, strength gains and potential improvements in endurance performance. A large portion of the strength exercises used in both the cycling^[17, 19, 28-31] and running literature^[12, 16] were open-chain, isolated and machine-based exercises (i.e. leg extension, seated hamstring curl, leg press, isometric plantar flexion). Stone & Stone^[50] states that strength programmes dominated by open-chain exercises may not provide adequate movement pattern specificity for optimal performance improvements in closed-chain sporting movements (i.e. running). As previously discussed, endurance sports require the hip, knee and ankle joint musculature to work in unison to produce force against the ground or pedal and provide locomotion. As a result of decreased mechanical specificity, the transferability of these strength exercises to performance may have been reduced. Although running can contain a combination of both open- and closed-chain movements, it is the closed-chain phase where force is produced against the ground to provide locomotion. Also, Stensdotter et al.^[52] demonstrated that there can be varying muscle activation patterns when an isolated, open-chain quadriceps exercise is compared to a multi-joint, closed-chain quadriceps exercise. These intra- and inter-muscular differences in exercises may complicate the learning and neural effects in the transfer of training process. Traditional multi-joint strength exercises, whether they are maximal- (i.e. squats, deadlifts & single-leg equivalents), explosive- (i.e. jump-squats, Olympic lift variations) or reactive-strength exercises (i.e. drop-jumps, sprints), are believed to be superior for eliciting optimal neuromuscular adaptations and increasing the force capabilities of the leg musculature.^[50] Future studies investigating the effect of strength training in endurance sports should programme these functionally superior exercises.

4.5.3 *Load and Velocity Prescription*

There are three main types of strength training: maximal-strength, explosive-strength (strength-speed & speed-strength) and reactive-strength training. Each can be categorised by velocity of the movement.^[38] All types of strength training were used in this review: reactive-,^[5, 13-15, 27] explosive-^[15] and maximal-strength orientated programmes.^[12, 16, 17, 19, 28-30, 31, 34, 36] Others used a mixed approach with no emphasis on a specific strength quality.^[17, 18, 35] A strength programme should be tailored to the current strength level of the athlete and evolves as they increase their force capabilities. Programming for a weak, or neuromuscular inefficient, athlete *can* be completely different (exercise, load, velocity, volume and frequency) to a strong athlete. Continual improvements in strong athletes require the development of programmes that target a specific strength quality (maximal-strength, strength-speed, speed-strength, and reactive-strength) in the force-velocity relationship.^[51] In contrast, athletes with low levels of strength, even though they may be a well-trained endurance athlete, can display improvements in neuromuscular function and force production from relatively non-specific and general strength programmes.^[53] This could be an explanation for why there were significant improvements in running economy from all three types of

THE EFFECT OF STRENGTH TRAINING ON PERFORMANCE IN ENDURANCE ATHLETES

strength training: reactive-,^[5, 13, 15] explosive-^[15] and maximal-strength interventions.^[12, 36] However, future studies that investigate longitudinal strength adaptations in endurance athletes should consider specifically prescribed programming for long-term gains.

Research in untrained subjects has shown that the neuromuscular adaptations from general strength training can result in a shift of the force–velocity curve in which force production is greater at any given velocity.^[54] Recent work from Cormie et al.^[53] found that in weak subjects, maximal-strength training not only improved the maximal force capabilities of the leg extensors, but the programme was also as effective as an explosive-strength programme in improving maximal power output. Further research from Dymond et al.^[55] found that subjects with higher levels of relative maximal-strength demonstrated superior reactive-strength ability. Dymond et al.'s^[55] work supports anecdotal evidence that reactive-strength, specifically the slow stretch-shortening cycle (i.e. a countermovement jump); can be improved in non-strength trained individuals following a period of maximal-strength training. In weak endurance athletes, especially where long-term improvements are the goal, a maximal-strength emphasised programme may initially be an efficient and effective training modality for improving several strength qualities together. Thus, weak endurance athletes may not necessarily need to place a focus on explosive- or reactive-strength training until a solid foundation of relative maximal-strength and neuromuscular efficiency is obtained. Nonetheless, reactive-strength can still be trained in low volume and supplemented alongside a maximal-strength orientated programme (i.e. basic plyometric progressions, stiff-leg pogos) and emphasis towards strength specificity can shift as the athlete enhances their neuromuscular ability.

4.5.4 The Interference Effect

As illustrated in Figure 1, appropriate strength training improves neuromuscular capacity, whereas endurance training targets both aerobic and anaerobic energy systems. However, recent molecular physiology research is starting to explain the intracellular signalling networks mediating exercise-induced skeletal muscle adaptations to both strength and endurance training stimuli. Simultaneously training for both strength and endurance may result in an acute compromised adaptation when compared with single-mode training^[56]. Strength training can activate the phosphatidylinositol 3-kinase (PI3-k)–Akt–mammalian target of rapamycin (mTOR) signalling pathway that regulates rate of protein synthesis, and over a prolonged period of time, muscle hypertrophy. Whereas endurance training activates another signalling cascade, the adenosine-monophosphate-activated protein kinase (AMPK)–p38 mitogen-activated protein kinase (MAPK)–peroxisome proliferator-activated receptor-gamma coactivator (PGC)-1 axis pathway. However, the activation of AMPK from the endurance training stimulus may interfere with, and inhibit, the mTOR signal for strength training-induced muscle protein synthesis^[56]. In short, an endurance-specific training session (i.e. LSD, tempo, interval) may inhibit the signalling pathway for optimal neuromuscular adaptation from the strength training stimulus. Nonetheless, molecular research in the area is in its infancy and there is much work to be undertaken before the information can be directly applied to the physical preparation of endurance athletes. Still, it is important that coaches are aware of the potential compromised adaptations when periodizing strength sessions in an endurance athlete's programme.

5. CONCLUSION & FUTURE DIRECTIONS

The present research available suggests the inclusion of strength training in an endurance athlete's programme for improved economy, muscle power and performance. It is important that future researchers and coaches are aware that muscular force-velocity adaptations are dependent upon the duration of the strength programme, the current strength-level of the athlete and the exercises administered (including the velocity and loads of the exercises). For long-term improvements in weak (neuromuscular inefficient) or non-strength trained endurance athletes, the present literature demonstrates that a general maximal-strength orientated programme may initially be the most appropriate and efficient method for improving maximal force, power and reactive-strength capabilities. Endurance athletes with high force capabilities may need to place a greater emphasis on specific explosive- and reactive-strength training to gain further improvements in performance. However, it is evident that further research is needed in this area. Future investigations should include valid strength assessments (i.e. squats, jump-squats, drop jumps) through a range of velocities (maximal strength ↔ strength-speed ↔ speed-strength ↔ reactive-strength), and administer appropriate programming (exercise, load & velocity prescription) over a long-term intervention period (> 6 months) for optimal transfer to performance.

Acknowledgements

The authors have no potential conflicts of interest that are directly relevant to the content of this review. This research is supported by funding from the University of Limerick Physical Education and Sport Science (PESS) Scholarship 2012.

REFERENCES

1. Bassett DR, Howley ET. Limiting factors for maximum oxygen uptake and determinants of endurance performance. *Med Sci Sports Exerc* 2000; 32(1): 70-84.
2. Paavolainen L, Nummela A, Rusko H. Muscle power factors and $\dot{V}O_{2max}$ as determinants of horizontal and uphill running performance. *Scand J Med Sci Sports* 2000; 10: 286-291.
3. Foster C, Lucia A. Running economy: the forgotten factor in elite performance. *Sports Med* 2007; 37(4-5): 316-9.
4. Fletcher JR, Esau SP, MacIntosh BR. Economy of running: beyond the measurement of oxygen uptake. *J Appl Physiol* 2009; 107: 1918-1922.
5. Paavolainen L, Hakkinen K., Hamalainen I, et al. Explosive strength training improves 5-km running time by improving running economy and muscle power. *J Appl Physiol* 1999; 86: 1527–1533.
6. Nummela A, Alberts M, Rijntjes RP, et al. Reliability and Validity of the Maximal Anaerobic Running Test. *Int J Sports Med* 1996; 17(2 Suppl.): 97 – 102.
7. McLaughlin JE, Howley ET, Bassett DR, et al. Test of the classic model for predicting endurance running performance. *Med Sci Sports Exerc* 2010; 42(5): 991-997.
8. Millet GP, Dréano P, Bentley DJ. Physiological characteristics of elite short- and long-distance triathletes. *Eur J Appl Physiol* 2003; 88: 427-430.

THE EFFECT OF STRENGTH TRAINING ON PERFORMANCE IN ENDURANCE ATHLETES

9. Zatsiorsky VM. *Science and Practice of Strength Training*. Champaign, IL, USA: Human Kinetics; 1995.
10. Karp JR. Training characteristics of qualifiers for the U.S. Olympic Marathon Trials. *Int J Sports Physiol Perform* 2007; 2: 72-92.
11. Maher CG, Sherrington C, Herbert RD, et al. Reliability of the PEDro scale for rating quality of randomized controlled trials. *Phys Ther* 2003; 83(8): 713–721.
12. Johnston RE, Quinn TJ, Kertzer R, et al. Strength training in female distance runners: Impact on running economy. *J Strength Cond Res* 1997; 11 (4): 224-229.
13. Spurrs RW, Murphy AJ, Watsford ML. The effect of plyometric training on distance running performance. *Eur J Appl Physiol* 2003; 89: 1-7.
14. Saunders PU, Telford RD, Pyne DB, et al. Short-term plyometric training improves running economy in highly trained middle and long distance runners. *J Strength Cond Res* 2006; 20(4): 947-954.
15. Berryman N, Maurel D, Bosquet L. Effect of plyometric vs. dynamic weight training on the energy cost of running. *J Strength Cond Res* 2010; 24(7): 1818-1825.
16. Fletcher JR, Esau SP, MacIntosh BR. Changes in tendon stiffness and running economy in highly trained distance runners. *Eur J Appl Physiol* 2010; 110: 1037-1046.
17. Jackson NP, Matthew SH, Reiser RF. High resistance / low repetition vs. low resistance / high repetition training: Effects on performance of trained cyclists. *J Strength Cond Res* 2007; 21(1): 289-295.
18. Levin GT, McGuigan MR, Laursen PB. (2009). Effect of concurrent resistance and endurance training on physiologic and performance parameters of well-trained endurance cyclists. *J Strength Cond Res* 2009; 23(8): 2280-2286.
19. Aagaard P, Andersen JL, Bennekou M, et al. Effects of resistance training on endurance capacity and muscle fiber composition in young cyclists. *Scand J Med Sci Sports* 2011; 21: 298 – 307.
20. Hoff J, Helgerud J, Wisløff U. Maximal strength training improves work economy in trained female cross-country skiers. *Med Sci Sports Exerc* 1999; 31(6): 870 – 877.
21. Hoff J, Gran A, Helgerud J. Maximal strength training improves aerobic performance. *Scand J Med Sci Sports* 2002; 12: 288-295.
22. Østerås H, Helgerud J, Hoff J. Maximal strength-training effects on force-velocity and force-power relationships explain increases in aerobic performance in humans. *Eur J Appl Physiol* 2002; 88: 255-263.
23. Millet GP, Jaouen B, Borrani F, et al. Effects of concurrent endurance and strength training on running economy and VO₂ kinetics. *Med Sci Sports Exerc* 2002; 34(8): 1351-9.
24. Hausswirth C, Argentin S, Bieuzen Y, et al. Endurance and strength training effects on physiological and muscular parameters during prolonged cycling. *J Electromyogr Kinesiol* 2009; 20: 330-339.
25. Bonacci J, Green D, Saunders PU, et al. Plyometric training as an intervention to correct altered neuromotor control during running after cycling in triathletes: A preliminary randomized controlled trial. *Phys Ther Sport* 2011; 12: 15-21.
26. Mikkola JS, Rusko HK, Nummela AT, et al. Concurrent endurance and explosive type strength training increases activation and fast force production of leg extensor muscles in endurance athletes. *J Strength Cond Res* 2007; 21(2): 613-620.

THE EFFECT OF STRENGTH TRAINING ON PERFORMANCE IN ENDURANCE ATHLETES

27. Mikkola J, Rusko H, Nummela A, et al. Concurrent endurance and explosive type strength training improves neuromuscular and anaerobic characteristics in young distance runners. *Int J Sports Med* 2007; 28(7): 602-611.
28. Rønnestad BR, Hansen EA, Raastad T. In-season strength maintenance training increases well-trained cyclists' performance. *Eur J Appl Physiol* 2010; 110 (6): 1269 – 1282.
29. Rønnestad BR, Hansen EA, Raastad T. Strength training affects tendon cross-sectional area and freely chosen cadence differently in non-cyclists and well-trained cyclists. *J Strength Cond Res* 2012; 26(1): 158-166.
30. Rønnestad BR, Hansen EA, Raastad T. Strength training improves 5-min all out performance following 185 min of cycling. *Scand J Med Sci Sports* 2011; 21(2): 250-259.
31. Rønnestad BR, Hansen EA, Raastad T. Effect of heavy strength training on thigh muscle cross-sectional area, performance determinants, and performance in well-trained cyclists. *Eur J Appl Physiol* 2010; 108: 965–975
32. Rønnestad BR, Kojedal Ø, Losnegard T, et al. Effect of heavy strength training on muscle thickness, strength, jump performance, and endurance performance in well-trained Nordic Combined athletes. *Eur J Appl Physiol* 2012; 112: 2241-2352.
33. Losnegard T, Mikkelsen K, Rønnestad BR, et al. The effect of heavy strength training on muscle mass and physical performance in elite cross country skiers. *Scand J Med Sci Sports* 2011; 21: 389-401.
34. Sunde A, Støren Ø, Bjerkaas M, et al. Maximal strength training improves cycling economy in competitive cyclists. *J Strength Cond Res* 2010; 24 (8): 2157-2165.
35. Bastiaans JJ, Van Diemen AB, Veneberg T, et al. The effects of replacing a portion of endurance training by explosive strength training on performance in trained cyclists. *Eur J Appl Physiol* 2001; 86: 79–84.
36. Støren O, Helgerud J, Støa EM, et al. Maximal strength training improves running economy in distance runners. *Med Sci Sports Exerc* 2008; 40: 1089– 1094.
37. Newton RU, Cormie P, Cardinale M. Principles of Athletic Testing. In: Cardinale M, Newton R, Nosaka K, editors. *Strength & Conditioning: Biological Principles and Practical Application*. Oxford, Wiley-Blackwell; 2011. pp. 255 – 270.
38. Siff M. *Supertraining*. 2nd ed. Denver, USA; 2003. p. 20.
39. Brughelli M, Cronin J. Influence of running velocity on vertical, leg and joint stiffness. Modelling and recommendations for future research. *Sports Med* 2008; 38(8): 647-667.
40. Schmidtbleicher D. Training for power events. In: Komi PV, editors. *The Encyclopedia of Sports Medicine*. Vol 3: Strength and Power in Sport. Oxford, UK: Blackwell; 1992. pp. 169-179.
41. Flanagan EP, Comyns TM. The use of contact time and the reactive strength index to optimize fast stretch-shortening cycle training. *Strength Cond J* 2008; 30(5): 32-38.
42. Jeukendrup AE, Craig NP, Hawley JA. The bioenergetics of world class cycling. *J Sci Med Sports* 2000; 3(4): 414-433.
43. Saunders PU, Pyne DB, Telford RD, et al. Factors affecting running economy in trained distance runners. *Sports Med* 2004; 34(7): 465-485.

THE EFFECT OF STRENGTH TRAINING ON PERFORMANCE IN ENDURANCE ATHLETES

44. Noakes TD. *Lore of Running*. 4th ed. Champaign, IL, USA: Human Kinetics; 2003. p 19-21.
45. Berg K. Endurance training and performance in runners. Research limitations and unanswered questions. *Sports Med* 2003; 33(1): 59-73.
46. Paavolainen LM, Nummela AT, Rusko HK. Neuromuscular characteristics and muscle power as determinants of 5 km running performance. *Med Sci Sports Exerc* 1999; 31: 124-130.
47. Noakes TD. Implications of exercise testing for prediction of athletic performance: a contemporary perspective. *Med Sci Sports Exerc* 1988; 20: 319 – 330.
48. Kraemer WJ, Ratamess NA. Fundamentals of resistance training: Progression and exercise prescription. *Med Sci Sports Exerc* 2004; 36(4): 674-688.
49. Appleby B, Newton RU, Cormie P. Changes in strength over a 2-year period in professional Rugby Union players. *J Strength Cond Res* 2012; 26 (9): 2538-2546.
50. Stone MH, Stone ME. Resistance training modes: a practical perspective. In: Cardinale, M., Newton, R. & Nosaka, K, editors. *Strength & Conditioning: Biological Principles and Practical Application*. Oxford: Wiley-Blackwell; 2011. p. 353.
51. Cormie P, McGuigan MR, Newton RU. Developing maximal neuromuscular power. Part 2 – training considerations for improving maximal power production. *Sports Med* 2011; 41(2): 125-146.
52. Stensdotter AK, Hodges PW, Mellor R, et al. Quadriceps activation in closed and in open kinetic chain exercise. *Med Sci Sports Exerc* 2003; 35(12): 2043-2047.
53. Cormie R, McGuigan MR, Newton RU. Adaptations in athletic performance after ballistic power versus strength training. *Med Sci Sports Exerc* 2010; 42(8): 1582-1598.
54. Kaneko M, Fuchimoto T, Toji H, et al. Training effect of different loads on the force-velocity relationship and mechanical power output in human muscle. *Scand J Med Sci Sports* 1983; 5(2): 50-55
55. Dymond C, Flanagan EP, Turner AP. The relationship between maximal strength and plyometric ability in Rugby players. *Rev Port Cein Desp* 2011; 11 (Supple. 2).
56. Hawley, J. Molecular responses to strength and endurance training: are they incompatible? *Appl Physiol Nutr Metab* 2009;34: 355 – 361.

THE EFFECT OF STRENGTH TRAINING ON PERFORMANCE IN ENDURANCE ATHLETES

SYSTEMATIC REVIEW

THE EFFECT OF STRENGTH TRAINING ON PERFORMANCE IN ENDURANCE ATHLETES

SYSTEMATIC REVIEW