

ULRR

Actor and partner effect of loneliness on episodic memory and verbal fluency: a dyadic multilevel analysis of romantic couples across 28 countries

Item Type	Article
Authors	Luchetti, Martina;Ledermann, Thomas;Aschwanden, Damaris;Nikitin, Jana;Ó Súilleabháin, Páraic;Stephan, Yannick;Terracciano, Antonio;Sutin, Angelina R
Citation	The Journals of Gerontology: Series B, Volume 77, Issue 12, December 2022, Pages 2202–2211
Publisher	Oxford University Press
Download date	2026-03-16 05:43:24
Item License	https://creativecommons.org/licenses/by-nc-sa/4.0/
Link to Item	https://doi.org/10.34961/researchrepository-ul.24592497

**Actor and Partner Effect of Loneliness on Episodic Memory and Verbal Fluency: A
Dyadic Multilevel Analysis of Romantic Couples across 28 Countries**

Martina Luchetti, Ph.D.,¹ Thomas Ledermann, Ph.D.,¹ Damaris Aschwanden, Ph.D.,¹ Jana Nikitin, Ph.D.,² Páraic S. O'Súilleabháin, Ph.D.,³ Yannick Stephan, Ph.D.,⁴ Antonio Terracciano, Ph.D.,¹ & Angelina R. Sutin, Ph.D.¹

¹ Florida State University, United States

² University of Vienna, Austria

³ University of Limerick, Ireland

⁴ University of Montpellier, France

Author Note

Correspondence concerning this article should be addressed to Martina Luchetti, Department of Behavioral Sciences and Social Medicine, Florida State University College of Medicine, 1115 West Call Street, Tallahassee, Florida 32306-4300. Email: martina.luchetti@med.fsu.edu

Abstract

Objectives: There is evidence that loneliness is detrimental for cognitive health. Most studies, however, do not consider the association between loneliness and cognition in the context of close relationships, such as a spouse or romantic partner. This study examines loneliness, experienced by both the individual and their romantic partner, and cognitive performance. **Methods:** Data were from 24,689 opposite-sex couples (49,378 participants) from 28 countries in the Survey of Health, Ageing and Retirement in Europe. Each couple participant reported on loneliness and completed memory and verbal fluency tasks. A multi-level sex-stratified analysis was used to account for the nested data structure and evaluate actor and partner effects of loneliness on cognitive performance for male and female partners. **Results:** Consistent with the literature, there were small actor effects of loneliness on memory and verbal fluency for both males and females: A person's own loneliness was associated negatively with their cognitive performance on both tasks. There were also small partner effects: A person with a partner who was lonely tended to have worse cognitive performance above and beyond their own loneliness. Actor and partners effects were similar for male and female partners, replicated in most countries, and generally held controlling for age, education, household size, and disease burden. For memory, loneliness effects were slightly stronger among older participants. **Discussion:** Both the experience of loneliness and loneliness of a partner have a negative association with cognitive health.

Keywords: actor loneliness, partner loneliness, cognitive function, romantic relationship, dyads

Loneliness is a concern for all age groups, including older adults (National Academies of Sciences, Engineering, and Medicine, 2020). Feeling lonely—which differs from being alone—is associated with increased risk for numerous health problems in middle and older adulthood (Hawkley & Cacioppo, 2010; National Academies of Sciences, Engineering, and Medicine, 2020), including cardiovascular problems (e.g., Valtorta et al., 2018), immune dysfunction (Shiovitz-Ezra & Parag, 2019), impaired sleep (Segrin & Burke, 2015), depression (Cacioppo et al., 2010), and even premature death (Holt-Lunstad et al., 2015). More recently, researchers have identified loneliness as one of the top risk factors for late-life dementia (Weiss et al., 2020; see also Lara, Martín-María, et al., 2019; Sundström et al., 2020; Sutin et al., 2020) and other cognitive impairments (Luchetti et al., 2020), independent of the effects of objective isolation (e.g., being unmarried or having few social contacts), health status, and other behavioral and generic risk factors for dementia (Luchetti et al., 2020; Sutin et al., 2020). Loneliness has also been associated with worse performance on tasks across multiple cognitive domains, including memory, verbal fluency, and backward digit span and a faster decline in general cognitive function over three years (Lara, Caballero, et al., 2019).

Studies that examined loneliness and cognition have only considered loneliness as an individual experience. Loneliness, however, is a phenomenon that occurs in relational contexts: It can spread from one individual to another (Cacioppo et al., 2009), particularly among close relationships, such a spouse or romantic partner (Ayalon et al., 2013; Stokes, 2017). Loneliness within spouses tends to be correlated and such feelings are associated with marital strain and lower perceived support (Ayalon et al., 2013; Ermer et al., 2020; Hsieh & Hawkley, 2018; Stokes, 2017). Hsieh and Hawkley (2018) found that individuals in ‘aversive’ marriages not only feel lonely themselves, but also contribute to their partners’ experience of loneliness. Individuals who tend to feel lonely may act negatively toward their

spouse, complain or criticize their partner, behavior that may ‘induce’ loneliness in the other person (Cacioppo et al., 2009; Hsieh & Hawkley, 2018). Individuals within a couple may also face similar circumstances that increase loneliness (i.e., the shared environment hypothesis), or tend to pair with a partner who shares similar levels of loneliness (i.e., the homophily hypothesis; see Cacioppo et al., 2009 and Ermer et al., 2020).

Independent of the mechanisms through which loneliness may spread within married or romantic partners, the experience of loneliness may have dyadic implications for health, and cognition in particular, among middle-aged and older couples. That is, in addition to one’s own level of loneliness, the loneliness of one’s partner may also have a negative association with cognitive health. There is some support for this hypothesis for health behaviors (e.g., sleep; Segrin & Burke, 2015) and cardiometabolic markers (e.g., HbA1c; Stokes & Barooah, 2021). Thus far, however, no study has examined the association between partner loneliness and cognitive health outcomes. Partner loneliness might be relevant for cognitive health for several reasons. First, a lonely partner may lead to more stressful interactions between spouses (Cacioppo et al., 2009; Hsieh & Hawkley, 2018), with consequences for both partners’ cognitive performance (Wilson et al., 2015). Second, a lonely partner may reduce the couple’s engagement in social activities and other cognitive stimulating activities, within and outside the household—activities that typically help maintain cognitive function with age (Krueger et al., 2009). Third, couples tend to share health behaviors that undermine health (Wilson et al., 2020) and that are also associated with loneliness (e.g., smoking, physical inactivity, etc., see Christiansen et al., 2016). Loneliness in one partner may impact cognition in the other partner through the shared, deleterious behaviors that result from feeling lonely.

The current study examines the cross-sectional association between loneliness and cognitive function using dyadic data from partnered couples in the Survey of Health, Ageing

and Retirement in Europe (SHARE). The aim of the study is to examine whether loneliness experienced both by the individual and their romantic partner is associated with cognitive performance. As illustrated in Figure 1, given the previous literature on loneliness and cognition, we hypothesize that the loneliness of Partner A will be associated with their own worse cognitive performance (*actor effect*). We further hypothesize that the loneliness of their partner, Partner B, will be associated negatively with their (Partner A's) cognitive performance (*partner effect*). We use a sex-stratified approach to examine possible differences in actor and partner loneliness associations among males and females, and explore whether these associations with cognition varied by age. We focus on memory and verbal fluency, as performance on these tasks is considered a clinically relevant marker for risk of subsequent cognitive impairment (Josefsson et al., 2019; Sutin et al., 2019).

Methods

Participants

This study used data from SHARE, a cross-national multi-disciplinary study of adults aged 50 and older and their spouses or romantic partners, regardless of age (Börsch-Supan, 2020, 2021). Data were drawn from 28 countries: France, Hungary, Portugal and Switzerland from Wave 4 (DOI: 10.6103/SHARE.w4.710), Austria, Belgium, Czech Republic, Denmark, Estonia, Germany, Israel, Luxemburg, the Netherlands, Sweden and Spain from Wave 5 (DOI: 10.6103/SHARE.w5.710), Croatia, Greece, Italy, and Slovenia from Wave 6 (DOI: 10.6103/SHARE.w6.710), and Bulgaria, Cyprus, Finland, Latvia, Lithuania, Malta, Poland, Romania, and Slovakia from Wave 8 (DOI: 10.6103/SHARE.w8.100). We selected these waves to maximize the number of couples and countries included in the analysis.¹ All participants in the analytic sample reported being married or in a committed relationship and lived with their partner at the time of the interview.² Across the waves, there were 49,378 participants (24,689 opposite-sex couples) who had complete data on loneliness, cognitive

function (memory and verbal fluency), age, education, and household size (i.e., the number of persons in the household). We excluded couples who had one or both partners with missing data on loneliness and/or cognition, age and/or education, and who reported discordant marital status at the time of the interview (see Supplemental Material, Figure S1, for a diagram of sample selection across waves).³ Because of the small number of same-sex couples in SHARE ($n = 37$), these couples were not included in the analysis.

SHARE is reviewed and approved by the Ethics Council of the Max Planck Society, Munich, Germany. Information on study design, sampling, and data availability can be found at: <http://www.share-project.org>.

Measures

Loneliness. Participants completed the 3-item version of the UCLA Loneliness Scale (Hughes et al., 2004; Mehrbrodt et al., 2019). Specifically, each member of the couple reported how much of the time they felt a lack of companionship, left out, and isolated from others on a 3-point scale (1 = often, 2 = some of the time, 3 = hardly ever or never). Responses were reverse-scored and the mean taken across the items, with higher scores reflecting greater loneliness (alpha reliabilities were .71 for males and .75 for females).

Cognitive Function. Participants completed standard measures of episodic memory and verbal fluency (Mehrbrodt et al., 2019). For the memory task, participants were read a list of 10 common words and asked to immediately recall the words. After a short delay in which other survey questions were answered, respondents were asked again to recall the 10 words. The score was the total number of words recalled across the immediate and delayed recall (range = 0–20); more words remembered indicated better memory function. For the verbal fluency task, participants were asked to name as many animals as possible in

60 seconds. The score was the total number of correct animals counted by the interviewer; more animals named indicated better processing speed and verbal abilities.

Covariates. Age in years, education, and household size (coded as 1= living with other relatives or children and 0 = living only with the partner) were selected as plausible confounders of the hypothesized associations (Hawkley et al., 2022). Education in SHARE is harmonized across countries using the 1997 International Standard Classification of Education with categories ranging from 0 (pre-primary level of education) to 6 (second stage of tertiary education). SHARE does not collect information on race/ethnicity. Additional analysis controlled for disease burden: Both partners reported whether they were ever diagnosed with the following diseases: heart attack, hypertension, stroke, lung disease, cancer (non-skin), arthritis, diabetes (score = 0-7). Sensitivity analysis further excluded participants who self-reported a diagnosis of Alzheimer's disease, dementia, or senility (yes/no). These additional covariates were included because experiencing a disease, either oneself or the disease of a spouse/partner, is related to both loneliness (Ferreira-Alves et al., 2014; Luchetti et al., 2021) and poor cognitive function (Comijs et al., 2009; Vitaliano et al., 2017).

Analytic Strategy

We first examined sex differences in loneliness and performance on memory and verbal fluency tasks using dependent-sample *t*-tests. Zero-order correlations were performed to test the associations between loneliness, cognitive performance, age, education, and health of each partner, and household size.

A two-level mixed model approach, with partners nested within countries, was used to estimate actor and partner effects (Kenny et al., 2006) and to account for the nonindependence within couples and countries. We conducted a sex-stratified analysis to

assess the association between loneliness and cognitive performance for male and female partners. We first predicted cognitive outcomes for males: Males' loneliness (actor loneliness) and females' loneliness (partner loneliness) were entered as predictors of males' performance on memory and verbal fluency, respectively (actor performance). We then used the same approach for females: Females' and males' loneliness were entered to predict females' performance on each task. We first estimated models with fixed effects of actor and partner loneliness on cognitive performance and then models with fixed and random effects. The models were then compared based on the Akaike Information Criterion (AIC). Using the best fitting model, we included the covariates: age of the actor, education of both partners, and household size. We did not include age of the partner because of the high correlation between the age of partners ($r = .89$). A sensitivity analysis excluded cases where one or both partners reported a diagnosis of Alzheimer's disease, dementia, or senility ($n = 438$). Additional analyses controlled for disease burden of each partner. Exploratory analyses tested whether the actor and partner effects varied by age. For the models with random effects, we allowed for different variances among the countries by using a variance components structure for the residuals. Effect sizes were calculated using the residual variances of the null and the full models. All analyses were performed using SPSS (version 25). All models applied the restricted maximum likelihood estimation.

Results

Descriptive Statistics and Correlations

Descriptive statistics and zero-order correlations for male and female partners are reported in Table 1 (see Supplemental Table S1 for statistics by country). There were a few noticeable differences between males and females (Cohen's $d > .10$): On average, females were lonelier (Cohen's $d = .11$) and had better performance on memory ($d = .22$) than

males. Males tended to be older ($d = .71$) and more educated ($d = .11$) compared to females. As expected, there were moderate correlations between males' and females' loneliness ($r = .35$) and between their cognitive performance ($r = .45$ for memory and $r = .54$ for verbal fluency).

Mixed Model Analysis

The AIC of the models with fixed effects predicting memory performance was 129,686 for males and 130,693 for females. The AIC for the models with fixed effects predicting verbal fluency was 165,793 for males and 164,990 for females. Adding the random effects of loneliness, AIC was 129,674 for males' and 130,678 for females' memory, and 165,757 for males' and 164,965 for females' verbal fluency. We therefore selected the models that included both random and fixed effects. Table 2 (Unadjusted) reports the actor and partner effects of loneliness on cognitive function. As expected, loneliness had a significant actor effect on memory and verbal fluency for both males and females: One's own loneliness was associated negatively with one's own cognitive performance. We also found evidence of significant partner effects that indicated that partner loneliness was related negatively to one's own memory and verbal abilities for both males and females. In other words, a person with a partner who was lonely tended to have poorer performance on both tasks above and beyond one's own loneliness. For memory, the size of the partner effects was about 40 percent of the size of the actor effects and for verbal fluency the partner effects were about half of the size of the actor effects. Across all models, the effects were small in size: f^2 was .02 for the models with loneliness indicating that actor and partner loneliness explained 2% of variance in cognitive performance relative to the unexplained variance of the null models without predictors. Actor and partner effects held accounting for age, education of both partners, and household size (see Table 2, Adjusted).

There was only one exception: Males' loneliness was no longer associated with females' memory when accounting for these covariates. The observed effects remained significant when further controlling for disease burden of both partners (see Supplemental Table S2) and excluding cases where one or both partners reported dementia or senility (see Table S3).⁴ Note that actor and partner effects of loneliness did not differ for males and females. Non-stratified analyses indicated non-significant interactions between actor loneliness and partner loneliness with sex, which indicated that the associations with cognitive performance were similar across males and females (see Table S4). The associations were also generally consistent across SHARE countries, with evidence of a significant partner effect observed in 18 out of the 28 individual countries and in the same direction in another 7 countries that generally had smaller samples (see Table S1).

Lastly, we tested whether the associations were moderated by age.⁵ Actor and partner effects on memory were slightly stronger for male partners at older ages: Experiencing loneliness and having a lonely partner were associated with worse memory, particularly in older males (actor loneliness \times actor age, $B = -.015$, $SE = .005$ [$\beta = -.016$], $p = .004$; partner loneliness \times actor age, $B = -.011$, $SE = .005$ [$\beta = -.012$], $p = .022$). For females, age moderated the actor effect but not the partner effect for memory: Loneliness was associated with worse memory, particularly for older females (actor loneliness \times actor age, $B = -.011$, $SE = .005$ [$\beta = -.012$], $p = .016$). No significant moderation emerged for the partner effect (partner loneliness \times actor age, $B = -.004$, $SE = .005$ [$\beta = -.004$], $p = .399$). Age did not moderate the association between loneliness and verbal fluency.

Discussion

This study examines the cross-sectional association between loneliness and cognitive function using dyadic data from partnered couples across 28 countries participating in the SHARE. There was evidence of small, significant actor and partner effects of loneliness on cognitive performance for both males and females. Consistent with the literature on loneliness and cognition (Lara, Caballero, et al., 2019), feeling lonely was associated negatively with a one's own performance on tasks of both memory and verbal fluency. To this literature, we add that participants with a partner who was lonely tended to have poorer performance on the cognitive tasks, above and beyond their own loneliness. Even if small, actor and partner effects generally replicated across countries, were roughly similar for male and female partners, and held when controlling for age, education, household size, and disease burden.

These results align with the accumulating evidence that feelings of loneliness are detrimental for cognitive health and increase risk of developing late-life dementia and other cognitive impairments (Lara, Martín-María, et al., 2019; Luchetti et al., 2020; Sutin et al., 2020). Previous studies have reported that loneliness is associated with poor cognitive performance in healthy adults (Kyröläinen & Kuperman, 2021; Lara, Caballero, et al., 2019), although the associations vary somewhat based on the cognitive domain and the psychological and sociodemographic variables included in the models (as noted by Boss et al., 2015). Our work supports a concurrent association between loneliness and worse word recall and verbal fluency, two tasks that are useful to identify individuals at risk for subsequent cognitive impairment (Josefsson et al., 2019; Sutin et al., 2019).

This work further expands the extant literature on loneliness and cognition by considering the context in which loneliness occurs, specifically the relational context with a

spouse or romantic partner. For many older adults, the relationship with their spouse or partner is the most important social connection and source of support. Within this relationship, the characteristics, behaviors, and experiences of each partner may have important implication for health (Choi et al., 2016; Stokes & Barooah, 2021), and in particular, for cognitive functioning. In line with this expectation, we found that feelings of loneliness, reported both by a participant and their partner were associated with poor memory recall and verbal fluency. That is, there was a partner as well as an actor effect of loneliness on cognition. Partner effects were about two-fifths in size of the actor effects for memory and about half of the actor effects for verbal functioning, which indicates a dyadic pattern in between of an 'actor-only pattern' (i.e., no partner effect) and a 'couple pattern' (i.e., equal actor and partner effect; Kenny & Ledermann, 2010). Of note, the effects were small in size: Actor and partner loneliness explained 2% of variance in cognitive performance. This is, however, not surprising and consistent with the literature (Boss et al., 2015; Lara, Caballero, et al., 2019). Cognitive health is, indeed, a complex and multi-determined phenomenon, with influences ranging from genetics to environmental factors (Tucker-Drob et al., 2013). Therefore, psychosocial and behavioral factors (including loneliness and other relational factors) are expected to explain only a small proportion of variance. Even if small, these effects are not negligible (Götz et al., 2022), especially when considering the documented effects of close relationships on health (Kiecolt-Glaser & Wilson, 2017; Wilson et al., 2020) and long-term consequences of loneliness on late-life cognition (Sundström et al., 2020; Sutin et al., 2020).

There are several potential mechanisms for the detrimental association between loneliness and cognitive function in the context of married or romantic couples. For example, loneliness can spread from one partner to the other (Ayalon et al., 2013; Stokes, 2017). Having a lonely partner may increase one's own level of loneliness and thereby impact

cognitive function. It is important to note, however, that partner loneliness was associated with cognitive performance above and beyond actor loneliness. There may thus be other mechanisms through which having a lonely partner may be associated with cognitive function. For instance, a lonely partner might criticize or withdraw affection and support from their spouse (Hsieh & Hawkley, 2018) that leads to stress and conflict within the couple, which in turn may detriment cognitive health (Wilson et al., 2015). A partner loneliness may be also an indicator of emotional disconnection in the dyad, which leads to negative emotions and psychological traits (i.e., neuroticism) that contribute to poor cognitive functioning (Foong et al., 2018; McHugh Power et al., 2020). Further, older couples tend to have similar behaviors that impact health and cognition (Wilson et al., 2020). The experience of loneliness by one partner might lead both members of the couple to disengage from social activities and interactions with others, with related consequences for cognitive function in daily life and over the long term (Wilson et al., 2015; Zhaoyang et al., 2021). Recent dyadic studies also found partner loneliness to be associated with health behaviors (Segrin & Burke, 2015) and physiological markers (Stokes & Barooah, 2021) that are associated with poor cognition. The present study found a partner effect of loneliness on cognitive function that, even if small, might have practical implications (Primbs et al., 2021). For example, when planning interventions directed to maintain cognitive functioning in older adulthood, researchers should consider the relational context in which a person lives.

It should be noted that, because of the correlational nature of our study, we cannot draw conclusions on causal pathways, nor the directionality of the associations. Longitudinal studies are needed to examine the co-evolution of loneliness within romantic partners and possible consequences on health and cognition (Ermer et al., 2020). Further, even though loneliness is a risk factor for cognitive decline and impairment decades later (Luchetti et al., 2020; Sundström et al., 2020), there are likely to be reciprocal associations of loneliness and

cognitive functioning (Zhong et al., 2017). For example, Zhong and colleagues (2017) found loneliness was associated with poor cognitive function at later assessments, while better cognitive function at the initial assessment was associated with decreases in loneliness over time. In the context of couples, a partner with better cognitive function may be more engaged and socially active, which may lower their own loneliness and also their partner's loneliness. This, in turn, could help support both partners' cognitive health. In contrast, when cognitive impairment is present in one of the partners, loneliness might increase, particularly for the spouse who transitions into the role of caregiver (Leggett et al., 2020; Luchetti et al., 2021).

Additional limitations and possibilities for future research are worth noting. First, our measure of sex was simply male and female. It is important to consider other variables related to cultural roles and relationship quality, in addition to sex, to examine whether aspects of the gendered experience and/or characteristics of the relationship operate in the partner effects. Second, although we found little moderating effect of age, there are other sociodemographic factors that were not available in SHARE (e.g., race/ethnicity) that merit attention in future studies. Third, couples with missing data were excluded from the analytic sample. Because of missingness, the results may underestimate the true associations, as noted in longitudinal studies on cognition (Salthouse, 2014). However, results were the same when imputing missing values for loneliness and cognition (see Supplemental Table S5). Fourth, this study aimed to test whether there was an association between partner loneliness with cognitive performance, in addition to the association with actor loneliness. The next step is to identify the mechanisms underlying these associations. That is, identifying possible mediators of loneliness and cognition, both at the individual level and within couples (e.g., affect or depression; McHugh Power et al., 2020). Finally, the social context and interactions outside the couple might be relevant to reduce loneliness and

protect cognition. In a recent ecological assessment study, Zhaoyang et al. (2021) found that having more daily social interactions (particularly pleasant interactions) was related to better cognitive performance on the same day and the following day. This type of micro-level analysis should be applied in future studies to understand contextual factors associated with loneliness, relationship dynamics, and their consequences on cognitive function and health.

In conclusion, results from this study illustrate the potential value of considering the relational context in which loneliness occurs when examining the association between loneliness and cognitive health. Loneliness is not the same as being alone and can be experienced even by married or partnered adults. Experiencing loneliness may not only have a negative association with one's own cognition but also the cognitive functioning of one's partner.

Accepted Manuscript

Acknowledgements

SHARE is coordinated by Dr. Axel Börsch-Supan at the Munich Center for the Economics of Aging, Max Planck Institute for Social Law and Social Policy. It is funded by the European Commission (Horizon 2020), the U.S. National Institute on Aging, and other national sources (see <http://www.share-project.org/organisation/funding.html>). SHARE data are available to user after registration at <http://www.share-project.org/data-access/share-conditions-of-use.html>. The current analyses were not pre-registered. Analytic syntaxes and outputs are available upon request to the corresponding author.

Funding Statement

Authors were supported by the National Institute on Aging of the National Institutes of Health under Award Numbers R01AG074573 and R01AG068093. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Conflicts of Interest

None.

Footnotes

¹ SHARE is an ongoing study. Since its start (Wave 1, 2004/06), new participants and new countries joined the study, with a total of 28 European countries and Israel participating in the last wave (Wave 8, 2019/20). In the current analyses, we included participants in a romantic relationship from across four waves: Wave 4 (i.e., first wave in which SHARE administered a 3-item measure of loneliness), Wave 5, Wave 6 and Wave 8. Wave 3 and Wave 7 were not considered because a life story interview was administered at these waves.

² Partners in SHARE are coupled through the variable 'coupleid'. This variable is provided only for individuals in a relationship and when both partners live in the same household. It is a system variable generated by SHARE based on the cover-screen information provided by one person within the household on behalf all other members. During the interview, each participant further responds to questions on their current marital or relationship status and whether changes occurred since the last interview. Because of the possible delay in time between the cover-screen and interview, there can be discordance between couple-ids and marital status, and marital status within a couple, in some cases. Participants who had a couple-id and a discordant marital status (i.e., either one or both partners reported to be separated, widowed or divorced at the time of the interview, $n = 1,354$) were exclude from the analysis.

³ Loneliness was measured as part of the drop-off questionnaire in Wave 4. This led to a higher number of couples with missing data on loneliness in Wave 4, as compared to the other waves in which loneliness was assessed as part of the interview (see Supplemental Material, Figure S1). Note that results were virtually the same when controlling for administration modality (loneliness as part of the interview vs. drop-off questionnaire) and when missing values for loneliness and cognition were imputed using multiple imputations (see Supplemental Table S5).

⁴ Because we aimed to identify an association between actor and partner loneliness with cognitive performance, we avoided overcorrection and inclusion of covariates that are potential mediators (e.g., depression) or colliders of the association. We conducted, however, additional analyses controlling for depressive symptoms. When controlling for actor depression, the effect of actor and partner loneliness on verbal fluency was reduced in size but remained significant (actor loneliness, $B = -.75$, $SE = .20$, $p = .001$ and partner loneliness,

$B = -.35$, $SE = .16$, $p = .033$ for males, and actor loneliness, $B = -.73$, $SE = .17$, $p < .001$ and partner loneliness, $B = -.41$, $SE = .16$, $p = .018$ for females). Actor but not partner loneliness was associated with memory controlling for depression in both males and females (actor loneliness, $B = -.33$, $SE = .07$, $p < .001$ for males and $B = -.37$, $SE = .06$, $p = p < .001$ for females). These analyses were exploratory and future work is needed to formally test mediation pathways between loneliness and cognition.

⁵ In additional exploratory analyses, we examined whether actor and partner effects of loneliness were further moderated by partners' disease burden and household size. For females, disease burden moderated some of the actor and partner effects. Specifically, partner loneliness was more detrimental for memory among females with better health (i.e., lower number of diseases; partner loneliness \times actor disease burden, $B = .121$, $SE = .048$, $p = .011$). Similarly, both actor and partner loneliness were more detrimental for verbal fluency among females with fewer diseases (actor loneliness \times actor disease burden, $B = .238$, $SE = .085$, $p = .005$; partner loneliness \times actor disease burden, $B = .217$, $SE = .099$, $p = .029$). There were no significant interactions between actor and partner loneliness and partner disease burden. Lastly, household size did not moderate actor and partner effects of loneliness on either memory or verbal fluency.

References

- Ayalon, L., Shiovitz-Ezra, S., & Palgi, Y. (2013). Associations of loneliness in older married men and women. *Aging & Mental Health, 17*(1), 33–39.
<https://doi.org/10.1080/13607863.2012.702725>
- Börsch-Supan, A. (2020). *Survey of Health, Ageing and Retirement in Europe (SHARE) Wave 7 (7.1.1)* [Data set]. SHARE–ERIC. <https://doi.org/10.6103/SHARE.W7.711>
- Börsch-Supan, A. (2021). *Survey of Health, Ageing and Retirement in Europe (SHARE) Wave 8 (1.0.0)* [Data set]. SHARE–ERIC. <https://doi.org/10.6103/SHARE.W8.100>
- Boss, L., Kang, D.-H., & Branson, S. (2015). Loneliness and cognitive function in the older adult: A systematic review. *International Psychogeriatrics, 27*(4), 541–553.
<https://doi.org/10.1017/S1041610214002749>
- Cacioppo, J. T., Fowler, J. H., & Christakis, N. A. (2009). Alone in the crowd: The structure and spread of loneliness in a large social network. *Journal of Personality and Social Psychology, 97*(6), 977–991. <https://doi.org/10.1037/a0016076>
- Cacioppo, J. T., Hawkley, L. C., & Thisted, R. A. (2010). Perceived social isolation makes me sad: 5-year cross-lagged analyses of loneliness and depressive symptomatology in the Chicago Health, Aging, and Social Relations Study. *Psychology and Aging, 25*(2), 453–463. <https://doi.org/10.1037/a0017216>
- Choi, H., Yorgason, J. B., & Johnson, D. R. (2016). Marital quality and health in middle and later adulthood: Dyadic associations. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences, 71*(1), 154–164.
<https://doi.org/10.1093/geronb/gbu222>
- Christiansen, J., Larsen, F. B., & Lasgaard, M. (2016). Do stress, health behavior, and sleep mediate the association between loneliness and adverse health conditions among older

people? *Social Science & Medicine*, 152, 80–86.

<https://doi.org/10.1016/j.socscimed.2016.01.020>

Comijs, H. C., Kriegsman, D. M. W., Dik, M. G., Deeg, D. J. H., Jonker, C., & Stalman, W.

A. B. (2009). Somatic chronic diseases and 6-year change in cognitive functioning among older persons. *Archives of Gerontology and Geriatrics*, 48(2), 191–196.

<https://doi.org/10.1016/j.archger.2008.01.005>

Ermer, A. E., Segel-Karpas, D., & Benson, J. J. (2020). Loneliness trajectories and correlates of social connections among older adult married couples. *Journal of Family*

Psychology, 34(8), 1014–1024. <https://doi.org/10.1037/fam0000652>

Ferreira-Alves, J., Magalhães, P., Viola, L., & Simoes, R. (2014). Loneliness in middle and old age: Demographics, perceived health, and social satisfaction as predictors.

Archives of Gerontology and Geriatrics, 59(3), 613–623.

<https://doi.org/10.1016/j.archger.2014.06.010>

Foong, H. F., Hamid, T. A., Ibrahim, R., & Haron, S. A. (2018). Mediating role of

neuroticism in the association between loneliness and cognitive function among community-dwelling older adults. *Australasian Journal on Ageing*, 37(4), 283–287.

<https://doi.org/10.1111/ajag.12555>

Götz, F. M., Gosling, S. D., & Rentfrow, P. J. (2022). Small effects: The indispensable

foundation for a cumulative psychological science. *Perspectives on Psychological Science*, 17(1), 205–215. <https://doi.org/10.1177/1745691620984483>

Hawkley, L. C., Buecker, S., Kaiser, T., & Luhmann, M. (2022). Loneliness from young

adulthood to old age: Explaining age differences in loneliness. *International Journal of Behavioral Development*, 46(1), 39–49. <https://doi.org/10.1177/0165025420971048>

- Hawkley, L. C., & Cacioppo, J. T. (2010). Loneliness matters: A theoretical and empirical review of consequences and mechanisms. *Annals of Behavioral Medicine*, *40*(2), 218–227. <https://doi.org/10.1007/s12160-010-9210-8>
- Holt-Lunstad, J., Smith, T. B., Baker, M., Harris, T., & Stephenson, D. (2015). Loneliness and social isolation as risk factors for mortality: A meta-analytic review. *Perspectives on Psychological Science*, *10*(2), 227–237. <https://doi.org/10.1177/1745691614568352>
- Hsieh, N., & Hawkley, L. (2018). Loneliness in the older adult marriage: Associations with dyadic aversion, indifference, and ambivalence. *Journal of Social and Personal Relationships*, *35*(10), 1319–1339. <https://doi.org/10.1177/0265407517712480>
- Hughes, M. E., Waite, L. J., Hawkley, L. C., & Cacioppo, J. T. (2004). A short scale for measuring loneliness in large surveys: Results from two population-based studies. *Research on Aging*, *26*(6), 655–672. <https://doi.org/10.1177/0164027504268574>
- Josefsson, M., Sundström, A., Pudas, S., Nordin Adolfsson, A., Nyberg, L., & Adolfsson, R. (2019). Memory profiles predict dementia over 23–28 years in normal but not successful aging. *International Psychogeriatrics*, 1–9. <https://doi.org/10.1017/S1041610219001844>
- Kenny, D. A., Kashy, D. A., & Cook, W. L. (2006). *Dyadic data analysis*. Guildford Press.
- Kenny, D. A., & Ledermann, T. (2010). Detecting, measuring, and testing dyadic patterns in the actor–partner interdependence model. *Journal of Family Psychology*, *24*(3), 359–366. <https://doi.org/10.1037/a0019651>
- Kiecolt-Glaser, J. K., & Wilson, S. J. (2017). Lovesick: How couples' relationships influence health. *Annual Review of Clinical Psychology*, *13*(1), 421–443. <https://doi.org/10.1146/annurev-clinpsy-032816-045111>

- Krueger, K. R., Wilson, R. S., Kamenetsky, J. M., Barnes, L. L., Bienias, J. L., & Bennett, D. A. (2009). Social engagement and cognitive function in old age. *Experimental Aging Research*, 35(1), 45–60. <https://doi.org/10.1080/03610730802545028>
- Kyröläinen, A.-J., & Kuperman, V. (2021). The effect of loneliness on cognitive functioning among healthy individuals in mid- and late-adulthood: Evidence from the Canadian Longitudinal Study on Aging (CLSA). *Frontiers in Psychology*, 12, 701305. <https://doi.org/10.3389/fpsyg.2021.701305>
- Lara, E., Caballero, F. F., Rico-Uribe, L. A., Olaya, B., Haro, J. M., Ayuso-Mateos, J. L., & Miret, M. (2019). Are loneliness and social isolation associated with cognitive decline? *International Journal of Geriatric Psychiatry*, 34(11), 1613–1622. <https://doi.org/10.1002/gps.5174>
- Lara, E., Martín-María, N., De la Torre-Luque, A., Koyanagi, A., Vancampfort, D., Izquierdo, A., & Miret, M. (2019). Does loneliness contribute to mild cognitive impairment and dementia? A systematic review and meta-analysis of longitudinal studies. *Ageing Research Reviews*, 52, 7–16. <https://doi.org/10.1016/j.arr.2019.03.002>
- Leggett, A. N., Choi, H., Chopik, W. J., Liu, H., & Gonzalez, R. (2020). Early cognitive decline and its impact on spouse's loneliness. *Research in Human Development*, 17(1), 78–93. <https://doi.org/10.1080/15427609.2020.1750293>
- Luchetti, M., Terracciano, A., Aschwanden, D., Lee, J. H., Stephan, Y., & Sutin, A. R. (2020). Loneliness is associated with risk of cognitive impairment in the Survey of Health, Ageing and Retirement in Europe. *International Journal of Geriatric Psychiatry*, 35(7), 794–801. <https://doi.org/10.1002/gps.5304>
- Luchetti, M., Terracciano, A., Stephan, Y., Aschwanden, D., & Sutin, A. R. (2021). Personality and psychological health in caregivers of older relatives: A case-control

study. *Aging & Mental Health*, 25(9), 1692–1700.

<https://doi.org/10.1080/13607863.2020.1758907>

McHugh Power, J., Tang, J., Kenny, R. A., Lawlor, B. A., & Kee, F. (2020). Mediating the relationship between loneliness and cognitive function: The role of depressive and anxiety symptoms. *Aging & Mental Health*, 24(7), 1071–1078.

<https://doi.org/10.1080/13607863.2019.1599816>

Mehrbrodt, T., Gruber, S., & Wagner, M. (2019). *SHARE: Scales and multi-item indicators*.

<http://www.share->

[project.org/fileadmin/pdf_documentation/SHARE_Scales_and_Multi-](http://www.share-project.org/fileadmin/pdf_documentation/SHARE_Scales_and_Multi-)

[Item_Indicators.pdf](http://www.share-project.org/fileadmin/pdf_documentation/SHARE_Scales_and_Multi-Item_Indicators.pdf)

National Academies of Sciences, Engineering, and Medicine. (2020). *Social isolation and loneliness in older adults: Opportunities for the health care system*. The National Academies Press.

Primbs, M., Pennington, C. R., Lakens, D., Silan, M. A. A., Lieck, D. S. N., Forscher, P. S.,

Buchanan, E. M., & Westwood, S. J. (2021). *Are small effects the indispensable foundation for a cumulative psychological science? A reply to Götz et al. (2022)*

[Preprint]. PsyArXiv. <https://doi.org/10.31234/osf.io/6s8bj>

Salthouse, T. A. (2014). Selectivity of attrition in longitudinal studies of cognitive

functioning. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences*, 69(4), 567–574. <https://doi.org/10.1093/geronb/gbt046>

Segrin, C., & Burke, T. J. (2015). Loneliness and sleep quality: Dyadic effects and stress effects. *Behavioral Sleep Medicine*, 13(3), 241–254.

<https://doi.org/10.1080/15402002.2013.860897>

- Shiovitz-Ezra, S., & Parag, O. (2019). Does loneliness 'get under the skin'? Associations of loneliness with subsequent change in inflammatory and metabolic markers. *Aging & Mental Health*, 23(10), 1358–1366. <https://doi.org/10.1080/13607863.2018.1488942>
- Stokes, J. E. (2017). Two-wave dyadic analysis of marital quality and loneliness in later life: Results from the Irish Longitudinal Study on Ageing. *Research on Aging*, 39(5), 635–656. <https://doi.org/10.1177/0164027515624224>
- Stokes, J. E., & Barooah, A. (2021). Dyadic loneliness and changes to HbA1c among older US couples: The role of marital support as stress buffer. *Journal of Aging and Health*, 33(9), 698–708. <https://doi.org/10.1177/08982643211006498>
- Sundström, A., Adolfsson, A. N., Nordin, M., & Adolfsson, R. (2020). Loneliness increases the risk of all-cause dementia and Alzheimer's Disease. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences*, 75(5), 919–926. <https://doi.org/10.1093/geronb/gbz139>
- Sutin, A. R., Stephan, Y., Luchetti, M., & Terracciano, A. (2020). Loneliness and risk of dementia. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences*, 75(7), 1414–1422. <https://doi.org/10.1093/geronb/gby112>
- Sutin, A. R., Stephan, Y., & Terracciano, A. (2019). Verbal fluency and risk of dementia. *International Journal of Geriatric Psychiatry*, 34(6), 863–867. <https://doi.org/10.1002/gps.5081>
- Tucker-Drob, E. M., Briley, D. A., & Harden, K. P. (2013). Genetic and environmental influences on cognition across development and context. *Current Directions in Psychological Science*, 22(5), 349–355. <https://doi.org/10.1177/0963721413485087>
- Valtorta, N. K., Kanaan, M., Gilbody, S., & Hanratty, B. (2018). Loneliness, social isolation and risk of cardiovascular disease in the English Longitudinal Study of Ageing.

European Journal of Preventive Cardiology, 25(13), 1387–1396.

<https://doi.org/10.1177/2047487318792696>

- Vitaliano, P. P., Ustundag, O., & Borson, S. (2017). Objective and subjective cognitive problems among caregivers and matched non-caregivers. *The Gerontologist*, 57(4), 637-647. <https://doi.org/10.1093/geront/gnv690>
- Weiss, J., Puterman, E., Prather, A. A., Ware, E. B., & Rehkopf, D. H. (2020). A data-driven prospective study of dementia among older adults in the United States. *PLOS ONE*, 15(10), e0239994. <https://doi.org/10.1371/journal.pone.0239994>
- Wilson, R. S., Boyle, P. A., James, B. D., Leurgans, S. E., Buchman, A. S., & Bennett, D. A. (2015). Negative social interactions and risk of mild cognitive impairment in old age. *Neuropsychology*, 29(4), 561–570. <https://doi.org/10.1037/neu0000154>
- Wilson, S. J., Peng, J., Andridge, R., Jaremka, L. M., Fagundes, C. P., Malarkey, W. B., Belury, M. A., & Kiecolt-Glaser, J. K. (2020). For better and worse? The roles of closeness, marital behavior, and age in spouses' cardiometabolic similarity. *Psychoneuroendocrinology*, 120, 104777. <https://doi.org/10.1016/j.psyneuen.2020.104777>
- Zhaoyang, R., Scott, S. B., Martire, L. M., & Sliwinski, M. J. (2021). Daily social interactions related to daily performance on mobile cognitive tests among older adults. *PLOS ONE*, 16(8), e0256583. <https://doi.org/10.1371/journal.pone.0256583>
- Zhong, B.-L., Chen, S.-L., Tu, X., & Conwell, Y. (2017). Loneliness and cognitive function in older adults: Findings from the Chinese Longitudinal Healthy Longevity Survey. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences*, 72(1), 120–128. <https://doi.org/10.1093/geronb/gbw037>

Table 1*Descriptive Statistics and Correlations among Study Variables*

	Males	Females	<i>t</i> -test	Correlations						
	<i>M</i> (<i>SD</i>)	<i>M</i> (<i>SD</i>)		1	2	3	4	5	6	7
1. Loneliness	1.21 (0.37)	1.26 (0.42)	-16.59**	.354**	-.124**	-.145**	.077**	-.053**	.106**	.004
2. Memory	8.84 (3.45)	9.65 (3.59)	-34.39**	-.150**	.452**	.448**	-.371**	.331**	-.169**	.072**
3. Verbal Fluency	19.79 (7.62)	20.07 (7.73)	-5.90**	-.164**	.491**	.538**	-.263**	.313**	-.097**	.009
4. Age (years)	66.32 (9.19)	63.22 (9.28)	111.56**	.055**	-.347**	-.241**	.888**	-.159**	.262**	-.336**
5. Education Level	2.95 (1.45)	2.81 (1.43)	17.06**	-.100**	.398**	.412**	-.212**	.592**	-.088**	-.013*
6. Disease Burden ^a	1.02 (1.04)	0.95 (1.03)	8.57**	.125**	-.194**	-.131**	.317**	-.154**	.241**	-.102**
7. Household Size (> 2)	28.4% (7,007)		--	.004	.059**	.003	-.373**	.007	-.125**	--

Note. *N* = 24,689 couples (49,378 individuals). Data were from SHARE Wave 4 (France = 1,135 couples, Hungary = 893, Portugal = 504, Switzerland = 909), Wave 5 (Austria = 1,036 couples, Belgium = 1,410, Czech Republic = 1,463, Denmark = 1,139, Estonia = 1,402, Germany = 1,677, Israel = 621, Luxemburg = 356, the Netherlands = 1,108, Sweden = 1,137, Spain = 2,010), Wave 6 (Croatia = 854 couples, Greece = 1,532, Italy = 1,613, Slovenia = 1,143), Wave 8 (Bulgaria = 262 couples, Cyprus = 155, Finland = 299, Latvia = 204, Lithuania = 363, Malta = 283, Poland = 397, Romania = 425, and Slovakia = 359). Correlations (Pearson's *r*s) for males and females are reported above and below the diagonal. For each variable, correlations between partners are in bold along the diagonal. Descriptive statistics and correlations for each country separately are reported in Table S1.

^a *N* size is slightly reduced due to missing values (*n* = 24,634 males and females).

* *p* < .05, ** *p* ≤ .01.

Table 2

Cognitive Task:	Memory						Verbal Fluency					
	Males			Females			Males			Females		
	<i>B</i>	<i>SE</i>	β	<i>B</i>	<i>SE</i>	β	<i>B</i>	<i>SE</i>	β	<i>B</i>	<i>SE</i>	β
Unadjusted												
Intercept	10.465**	0.174		11.276**	0.205		22.794**	0.681		23.049**	0.763	
<i>Loneliness</i>												
Actor	-0.998**	0.080	-	-0.986**	0.062	-				-1.798**	0.176	-
Loneliness			.11				12	-1.866**	0.227	.09		.10
Partner	-0.394**	0.072	-	-0.386**	0.094	-				-0.884**	0.189	-
Loneliness			.05				.04	-0.864**	0.177	.05		.04
<i>Estimate</i>												
<i>Variance</i>												
Residual	11.118**	0.100		11.576**	0.104		47.835**	0.431		46.326**	0.418	
Intercept	0.589**	0.199		0.905**	0.283		11.750**	3.466		15.151**	4.364	
Actor Slope	0.058	0.039		0.014	0.024		0.846*	0.398		0.431*	0.221	
Partner Slope	0.047	0.035		0.115*	0.059		0.430	0.226		0.449	0.278	
Adjusted												
Intercept	15.718**	0.221		15.337**	0.234		29.738**	0.707		28.005**	0.760	
<i>Loneliness</i>												
Actor	-0.712**	0.068	-	-0.699**	0.054	-	-1.413**	0.211	-	-1.271**	0.162	-
Loneliness			.08				.08		.07			.07
Partner	-0.188**	0.061	-	-0.151	0.077	-	-0.494**	0.153	-	-0.497**	0.166	-
Loneliness			.02				.02		.03			.02
<i>Covariates</i>												
Actor Age	-0.119**	0.002	-	-0.110**	0.002	-	-0.176**	0.005	-	-0.166**	0.005	.20
			.32				.28		.21			
Actor Education	0.495**	0.017	.21	0.630**	0.018	.25	0.796**	0.037	.15	1.339**	0.037	.25
Partner Education	0.212**	0.017	.09	0.176**	0.017	.07	0.502**	0.038	.09	0.220**	0.035	.04
Household Size	-0.198**	0.046	-	-0.176**	0.048	-	-0.192	0.100	-	-0.225*	0.098	-
			.06				.05		.02			.03
<i>Estimate</i>												
<i>Variance</i>												
Residual	8.983**	0.081		9.262**	0.083		42.152**	0.380		39.187**	0.353	
Intercept	0.325**	0.112		0.522**	0.166		8.968**	2.664		11.797**	3.411	
Actor Slope	0.033	0.026		0.007	0.019		0.721*	0.338		0.369*	0.176	
Partner Slope	0.027	0.024		0.063	0.038		0.267	0.160		0.317	0.209	

Actor and Partner Effects of Loneliness on Memory and Verbal Fluency

Note. $N = 24,689$ couples ($49,378$ individuals). Unadjusted reports the models without covariates; Adjusted reports the models with basic covariates. Because we did not test cross-level interactions, individual-level variables were not centered. SPSS produces only unstandardized coefficients in multi-level modeling. Therefore, we calculated standardized coefficients using the formula: β standardized = (B unstandardized \times standard deviation explanatory variable)/standard deviation outcome variable; β s are reported for actor and partner effects of loneliness and covariates.

* $p \leq .05$, ** $p \leq .01$.


Accepted Manuscript

Figure 1

Model of the Hypothesized Actor and Partner Effects

Accepted Manuscript

Figure 1



Accepted Man